

Vaccine	Trade Name	Where to store	Acceptable Temperature Range	Diluent Storage	Specific Expiration after Opened/Reconstituted	Protect from Light	Other Comments
Adenovirus	Adenovirus Type 4 and Type 7	Refrigerator	2°C to 8°C (36°F to 46°F)		May be used until expiration date.		Keep bottles tightly closed and protect from moisture. Do NOT remove desiccant canister from bottles.
	BioThrax®	Refrigerator	2°C to 8°C (36°F to 46°F)		Once the stopper of the multi-dose vial has been pierced, the vial must be discarded within 28 days.		Shake well before use.
Cholera	Vaxchora®	Refrigerator	2°C to 8°C (36°F to 46°F)	Pour 100 mL of cold or room temperature (41°F-72°F; 5°C-22°C) purified bottled or spring bottled water into a clean, disposable cup. Do not use tap water, sparkling (carbonated) water, non-purified or non-spring bottled water, other beverages, or other liquids.	Must be consumed within 15 minutes of reconstitution. If the packets are reconstituted in the improper order, the vaccine must be discarded.	Yes	Protect from light and moisture. Reconstitution should be completed within 15 minutes of removing from the refrigerator. When out of refrigerated storage, packets should not be exposed to temperatures above 80°F (27°C).
	Daptacel® Infanrix® Kinrix®						Shake well before use.
All DTap vaccines	Pediarix®	Refrigerator	2°C to 8°C (36°F to 46°F)				Shake well before use.
	Pentacel®						Shake well before use.
Hepatitis A	Quadracel® Vaxelis®	Refrigerator	2°C to 8°C (36°F to 46°F)	Yes – store in refrigerator	Use immediately after reconstitution.		Shake well before use.
	Havrix®						Shake well before use.
	VAQTA®						Shake well before use.
	Engerix-B®						Shake well before use.
Hepatitis B	Heplisav-B® Recombinax HB®	Refrigerator	2°C to 8°C (36°F to 46°F)				Shake well before use.
	Twinnrix®	Refrigerator	2°C to 8°C (36°F to 46°F)			Yes	Shake well before use.
Hib vaccines	ActHIB®	Refrigerator	2°C to 8°C (36°F to 46°F)	Yes – store in refrigerator	Use within 24 hours of reconstitution.		Shake well before use. If the vaccine is not administered immediately, shake the solution well again before administration.
	Hiberix® PedvaxHIB®	Refrigerator	2°C to 8°C (36°F to 46°F)	Yes – store in refrigerator or at room temperature	Use within 24 hours of reconstitution.	Yes	If the vaccine is not administered immediately, shake the solution well again before administration. Shake well before use.
HPV	Gardasil 9®	Refrigerator	2°C to 8°C (36°F to 46°F)			Yes	Shake well before use. Administer as soon as possible after being removed from refrigeration.

Continued on Next Page