

DoD Opioid Overdose Education and Naloxone Distribution (OEND) Program: A Guide for Prescribing Naloxone **XX Month 20XX**



Pre-Training Assessment



- Online participants can access the pre-training survey by visiting:
- [\[Insert Link\]](#)

Learning Objectives



- Understand Department of Defense (DoD) Opioid Overdose Education and Naloxone Distribution (OEND) Program:
 - ❑ Use DoD OEND's **Quick Reference Guide** to understand when and how to prescribe naloxone to your patients
 - ❑ Use **CarePoint** to look up Risk Index for Overdose or Serious Opioid-Induced Respiratory Depression (RIOSORD) scores and Morphine Equivalent Daily Dose (MEDD)
 - ❑ Know what **key messages** to use when discussing naloxone with your patients

Agenda



■ Naloxone Overview

- OEND Program
- Policy vs. Active Implementation

■ Review the Quick Reference Guide

- Assess/Offer (CarePoint)
- RIOSORD and MEDD Assessment Tools (Case Example)
- Notify & Educate
- Document

■ Conclusion

- Key Takeaways
- References
- Questions

Discussion Touch Point



- Have you prescribed naloxone in the past?
 - What prompted you to prescribe naloxone?
- Describe your experiences with prescribing naloxone.
 - What are some concerns you have about co-prescribing naloxone to your patients?

OEND Program

- **Mission:** Reduce opioid-related overdoses and deaths
- **Goal:** Increase co-prescribing of naloxone

■ Policy Alignment:

- ❑ DoD/VA CPG for Opioid Therapy for Chronic Pain ([link](#))
- ❑ DHA-PI 6025.04: Pain Management and Opioid Safety in the Military Health System (MHS) ([link](#))
- ❑ DHA-PI 6025.07: Naloxone Prescribing and Dispensing by Pharmacists in Military Treatment Facilities (MTFs) ([link](#))



Nasal Spray
Narcan

Policy vs. Active Implementation




Figure 1: Percent of Patients with a RIOSORD > 32 Dispensed an Opioid Who Were Prescribed Naloxone in the Past Year

“Medically Ready Force...Ready Medical Force”

Review the Quick Reference Guide



- Developed to help prescribers and pharmacists understand when and how to prescribe naloxone
- Risk Criteria
 - RIOSORD>32
 - MEDD>50
 - Opioid/Benzodiazepine combination use
 - On long-term opioid therapy

When to Prescribe Naloxone?
A Quick Reference Guide for Military Health System Providers

Assess/Offer

Prescribe naloxone if a patient

- Has a Morphine Equivalent Daily Dose (MEDD) > 50
- Has a Risk Index for Overdose or Serious Opioid-Induced Respiratory Depression (RIOSORD) Score > 32
 - o RIOSORD scores located on the Opioid Registry and Patient Look up Tool at <https://carepoint.health.mil/>
 - o New diagnoses and prescriptions impact RIOSORD scores, but scores are not re-calculated in real-time. Manual calculations are recommended (see page 2 for worksheet).
- Check prescription drug monitoring program (PDMP) database to assess outside opioid prescriptions: <https://www.health.mil/Military-Health-Topics/Access-Cost-Quality-and-Safety/Access-to-Healthcare/Pharmacy-Program/Prescription-Drug-Monitoring-Program-Procedures>
- Has a benzodiazepine co-prescription with opioids
- Is on long-term opioid therapy
- Is considered at risk per your clinical judgment
- Directly requests naloxone

Notify

- The patient's other providers of opioid and naloxone prescriptions

Notify & Educate

Educate Patients...

Review Brochures

- "Save a Life: How to Prevent Opioid Poisoning Deaths" A Quick Reference Guide for Patients and Caregivers

Safeguard

- Properly dispose of unused opioids to protect others
- Know where naloxone is stored in the event of an overdose
- Involve family member(s) in education, if possible
- Recognize signs and symptoms of an overdose

Respond to an overdose

- Administer naloxone
- Contact 911 immediately

Document

Document in the electronic health record

- If naloxone was prescribed and why (e.g., risk indicated, patient request, clinical judgment)
- If naloxone was offered, but declined
- RIOSORD score
- High-risk opioid alerts and risk assessment results (including urine drug test results, PDMP findings)

Published by the Defense & Veterans Center for Integrative Pain Management (DVCIPM) in collaboration with DHA's Research and Development Directorate

U.1

Genetics and pharmacokinetics cannot account for all individual differences in genetics and pharmacokinetics.

Accessing CarePoint, MHSPHP, and the DHA Opioid Registry



- A one-pager with step-by-step instructions will be emailed to participants.
- Look up MHS-direct care RIOSORD scores, MEDD calculations, and opioid-related prescriptions (Rxs) on CarePoint: <https://carepoint.health.mil>



Instructions to Access Defense Health Agency (DHA) Opioid Registry

Naloxone and opioid-related information is displayed in the DHA Opioid Registry, located on CarePoint. Below are instructions to access the Opioid Registry on CarePoint to assist you in your naloxone assessment.

How to Access CarePoint

1. Using a computer with a CAC reader and internet access, go to <https://carepoint.health.mil>
2. Click the "Continue" button.
3. Use your CAC E-mail certificate and enter your PIN number to access the site.

How to go to the Military Health Service Population Health Portal (MHSPHP)

4. From the CarePoint home page, click on the "Apps" dropdown at the top menu bar, then select "All Apps."
5. Scroll down the list of Apps to the "MHSPHP" App icon.
6. Click on "Favorite" under the description to display it on your CarePoint home page.
7. Click on the MHSPHP icon to enter the App.

How to request Clinical Registries on Carepoint Protected Health Information (PHI) access on MHSPHP

8. Click on the red "Request PHI Access" link in the top left corner.
(Or go to: <https://carepoint.health.mil/sites/mhsphp/Site/Pages/RequestAccess.aspx>)
 - a. If you are a new Composite Health Care System (CHCS) user or a user that has a new location. Go to #9.
 - b. If you do not have a CHCS account. Go to #13.

CHCS Users (new user or new location) requesting PHPM access:

9. Click on the "Validate CHCS Account" button.
(Or go to the following link: <https://carepoint.health.mil/siteassets/PHI/PHIValidation.aspx>)
10. Type in the Military Treatment Facility (MTF) name or Defense Medical Information System of your MTF in the "Select MTF Box" and it will autofill with options.
11. Enter in your CHCS username and password. Click "Validate" when finished. (New passwords have a 24-hour delay prior to validation.)
12. If your account has been validated, click on the "Refresh" button and you can access the PHPM registries located on the left side under the "Clinical Registries" drop down menu. Go to #15.

Non-CHCS Users requesting PHPM access

13. Download the "Request Access Form" by either clicking the "Request Access" button or going to: <https://carepoint.health.mil/sites/mhsphp/PHPM%20Linked%20Documents/MHSPHPAccessRequest.pdf>
14. Complete the form electronically and send to the identified Service POCs listed on the webpage.

How to Access DHA Opioid Registry (with MHSPHP access)

15. Click the "Clinical Registries" on the left side navigation menu, then select "Opioid Management."
16. To set a preferred layout view of the data columns, click the icon at the top of the registry:
 - a. Manually customize the columns by selecting "Rearrange" tab.
 - b. Access public displays and filters by selecting "My Layouts" tab and clicking apply (ex.: "Simplified Registry")
17. To customize filters, click the button at the top of the registry:
 - a. On the "Filters" tab, select your own filters (ex.: RIOSORD greater than 32)
 - b. Select pre-made and custom-made filters on the "My Filters" tab (ex.: On long term opioid therapy with appointments today).

****Reminder to also access your state's local Prescription Drug Monitoring Program (PDMP)****


Look up your state's local PDMP: <https://www.health.mil/Military-Health-Topics/Access-Cost-Quality-and-Safety/Access-to-Healthcare/Pharmacy-Program/Prescription-Drug-Monitoring-Program-Procedures>

Created by the Defense & Veterans Center for Integrative Pain Management (DVCIPM) in collaboration with DHA J-9 Research Practice Integration
Last Updated June 2020

Instructions to Access DHA Opioid Registry

"Medically Ready Force...Ready Medical Force"

TUTORIAL



“Medically Ready Force...Ready Medical Force”

Quick Reference Guide: RIOSORD and MEDD Assessment Tools



■ Option to manually calculate two key indicators for whether your patient should be prescribed naloxone:

- RIOSORD Score > 32
- MEDD > 50

Risk Index for Overdose or Serious Opioid-Induced Respiratory Depression (RIOSORD): Calculate risk by completing RIOSORD assessment		Circle if "YES"
In the past 6 months, has the patient had a health care visit (outpatient, inpatient, or ED) involving:		
• Opioid dependence?		15
• Chronic hepatitis or cirrhosis?		9
• Bipolar disorder or schizophrenia?		7
• Chronic pulmonary disease? (e.g., emphysema, chronic bronchitis, asthma, pneumoconiosis, asbestosis)		5
• Chronic kidney disease with clinically significant renal impairment?		5
• Active traumatic injury, excluding burns? (fracture, dislocation, contusion, laceration, wound)		4
• Sleep apnea?		3
Does the patient consume:		
• Extended release or long acting (ER/LA) formulation: An ER/LA formulation of any prescription opioid or opioid with long and/or variable half-life? (e.g., OxyContin, Oramorph-SR, methadone, fentanyl patch, levorphanol)		9
• Methadone? (Methadone is a long-acting opioid, so also circle for "ER/LA formulation")		9
• Oxycodone? (If it has an ER/LA formulation [OxyContin], also circle for "ER/LA formulation")		3
• A prescription antidepressant? (e.g., fluoxetine, citalopram, venlafaxine, amitriptyline)		7
• A prescription benzodiazepine? (e.g., diazepam, alprazolam)		4
Is the patient's current maximum prescribed opioid dose: (Use Opioid Daily Dose Conversion Table below)		
• ≥100 mg morphine equivalents per day?		16
• 50 – <100 mg morphine equivalents per day?		9
• 20 – <50 mg morphine equivalents per day?		5
In the past 6 months, has the patient:		
• Had 1 or more ED visits?		11
• Been hospitalized for 1 or more days?		8
TOTAL SCORE (add up "YES" response values).		If score > 32, PRESCRIBE NALOXONE →
Opioid Daily Dose Conversion Table:		
Calculate Morphine Equivalent Daily Dose (MEDD) by multiplying daily dose for each opioid by the conversion *		
Type of Opioid (doses in mg/day except where noted)	Conversion Factor	MEDD (enter calculation here)
• Buprenorphine patch	12.6	
• Buprenorphine tab or film	10	
• Butorphanol (Stadol)	7	
• Codeine	0.15	
• Fentanyl transdermal (in mcg/hr)	2.4	
• Hydrocodone	1	
• Hydromorphone	4	
• Meperidine	0.1	
• Methadone		
○ 1-20 mg/day	4	
○ 21-40 mg/day	8	
○ 41-60 mg/day	10	
○ ≥ 61-80 mg/day	12	
• Morphine	1	
• Oxycodone	1.5	
• Oxymorphone	3	
• Tapentadol IR	0.4	
• Tramadol	0.1	
<small>*These dose conversions are estimated and cannot account for all individual differences in genetics and pharmacokinetics.</small>		

CASE STUDY



“Medically Ready Force...Ready Medical Force”

Case Example

Patient John Smith comes in for an outpatient visit regarding his chronic lower back pain.

He has been on hydrocodone for about a year. He requests a renewal of his prescription.

You have not yet established an account in CarePoint but want to determine if you should prescribe naloxone.

He currently has a prescription for citalopram for mild depression.

His current average daily opioid dosage is 52 mg morphine equivalent dosage per day.

The patient had one emergency department visit 4 months ago and was hospitalized for 3 days.

Based on the information provided, should you prescribe naloxone?

Risk Index for Overdose or Serious Opioid-Induced Respiratory Depression (RIOSORD): Calculate risk by completing RIOSORD assessment	Circle if "YES"
In the past 6 months, has the patient had a health care visit (outpatient, inpatient, or ED) involving:	
• Opioid dependence?	15
• Chronic hepatitis or cirrhosis?	9
• Bipolar disorder or schizophrenia?	7
• Chronic pulmonary disease? (e.g., emphysema, chronic bronchitis, asthma, pneumoconiosis, asbestosis)	5
• Chronic kidney disease with clinically significant renal impairment?	5
• Active traumatic injury, excluding burns? (fracture, dislocation, contusion, laceration, wound)	4
• Sleep apnea?	3
Does the patient consume:	
• <u>Extended release or long acting (ER/LA) formulation</u> : An ER/LA formulation of any prescription opioid or opioid with long and/or variable half-life? (e.g., OxyContin, Oramorph-SR, methadone, fentanyl patch, levorphanol)	9
• Methadone? (Methadone is a long-acting opioid, so also circle for "ER/LA formulation")	9
• Oxycodone? (If it has an ER/LA formulation [OxyContin], also circle for "ER/LA formulation")	3
• A prescription antidepressant? (e.g., fluoxetine, citalopram, venlafaxine, amitriptyline)	7
• A prescription benzodiazepine? (e.g., diazepam, alprazolam)	4
Is the patient's current maximum prescribed opioid dose: (Use Opioid Daily Dose Conversion Table below)	
• ≥ 100 mg morphine equivalents per day?	16
• 50 – <100 mg morphine equivalents per day?	9
• 20 – <50 mg morphine equivalents per day?	5
In the past 6 months, has the patient:	
• Had 1 or more ED visits?	11
• Been hospitalized for 1 or more days?	8
TOTAL SCORE (add up "YES" response values).	If score > 32, PRESCRIBE NALOXONE → 35

Quick Reference Guide: Notify & Educate

DHA-PI 6025.07 includes a brochure on opioid safety and naloxone administration that you can distribute to patients and their caregivers. This brochure is currently being finalized and will soon be available online at: health.mil/opioidsafety

SAVE A LIFE: HOW TO PREVENT OPIOID POISONING DEATHS

A Quick Reference Guide for Patients and Caregivers

What are opioids?
Opioids are drugs that may be used in pain management. Everyone who takes opioids is at risk for opioid poisoning (overdose), even if taken as prescribed, and should take appropriate precautions.

Common Opioids Include:

GENERIC NAME	BRAND NAME
Hydrocodone	Vicodin, Lortab, Lotab, Norco, Zilbyte
Oxycodone	Percocet, OxyContin, Roxicodone, Percodan
Morphine	MSContin, Kadian, Erbelex, Avanza
Codaine	Tylenol with Codaine, Tyco, Tylenol #3
Fentanyl	Duragesic, Actiq
Hydroxycodone	Dilaudid
Oxycodone	Opiana
Meprobamate	Demerol
Methadone	Dolophine, Methadone
Buprenorphine	Suboxone, Subutex, Zubsolv, Bunavail, Butrans

Heroin is also an opioid.

OPIOID SAFETY TIPS

DO take opioid and non-opioid medications as prescribed

DO inform all providers that you are taking opioids

- Tell your primary provider if another provider prescribes an opioid for you

DO be cautious about driving or operating machinery, especially if you feel sleepy or drowsy

DO get help from family and friends

- Tell them that you take opioids
- Ask them to help you take opioids safely
- Tell them where you keep the naloxone kit and how to use it

DON'T take extra doses of opioids

- You could overdose and die

DON'T drink alcohol or use "street" drugs when taking opioids; they can impair your ability to use opioids safely and can cause severe harm or death

DON'T share, give away, or sell your opioids

- This is dangerous and illegal!

If you've been taking opioids daily for more than a few weeks, DON'T stop taking opioids on your own


- You may feel flu-like withdrawal symptoms
- Your provider can help you stop safely
- You may overdose if you start taking opioids again after an opioid-free break

ADDITIONAL RESOURCES
Local Emergency Services: 911
For other patient and caregiver resources, please visit: health.mil/opioidsafety




IN CASE OF OPIOID POISONING (OVERDOSE)


What is naloxone?
Naloxone (brand name: Narcan) is a temporary antidote for an opioid overdose. Should an overdose occur, naloxone will temporarily restore your breathing. If you are prescribed opioids, your doctor or pharmacist may talk to you about or prescribe naloxone. If you or someone you know is taking opioids, you can request naloxone.
Someone else has to administer this medication to the person experiencing an overdose. Be sure to tell your family members and friends where you keep your naloxone, and teach them how to use it.
If you use naloxone, follow up with your provider.




- Check responsiveness**
Look for any of the following:

 - No response even if you shake them, say their name, or do a sternal rub
 - Breathing slows or stops
 - Lips and fingernails turn blue or gray
 - Skin gets pale or clammy
- Call 911 and give naloxone**
If no reaction in 2-3 minutes, give second naloxone dose in the other nostril. (medication comes in two packs)
This nasal spray needs no assembly and can be sprayed up one nostril by pushing the plunger.


Do not test device - each device can only be used once.


- Follow 911 dispatcher instructions**
Dispatcher may provide instructions for rescue breathing and/or CPR.




>> Stay with person until help arrives

For video instructions, use your phone's camera to scan the QR code



For more information about accessing naloxone, talk to your pharmacist or provider.



How to Talk to Patients and Caregivers about Naloxone

■ *Example Scenario (Role Play)*

■ Key messages/talking points:

- Opioid use disorder is a pain management issue
- Naloxone is a lifesaving precaution; it does not lead to increased drug abuse
- When managing pain, use opioids as prescribed



[Video](#) – How to use the VA Naloxone Nasal Spray (from the Veterans Health Administration)

ROLE PLAY



“Medically Ready Force...Ready Medical Force”

Scenario 1

I don't need medication to prevent overdose. I have been taking the medication for a long time and I don't have any problems with it.

Patient

What would you say?

Clinician

Are you saying the medication that I was prescribed is dangerous?

Patient

What would you say?

Clinician

Scenario 2

Are you saying you think I abuse drugs? I'm not a drug addict!

Patient

What would you say?

Clinician



Scenario 3

Ok, I'll think about it, but no thanks I don't want to take the prescription with me today.

Patient

What would you say?

Clinician



Scenario 4

How do I use naloxone?

Patient

What if I am unconscious and cannot administer naloxone myself?

Patient

What would you say?

Clinician

What would you say?

Clinician

Discussion Touch Point



- What did the pharmacist do that was useful in addressing the patient's concerns?
- What strategies would you have used to address them?
- Have you discussed naloxone with any of your patients?
 - Were they receptive?
 - What were some barriers you encountered?

TRIVIA



“Medically Ready Force...Ready Medical Force”

Question 1



True or False?

- My patient does not have an addiction problem, so they are not at risk for an opioid overdose.

Question 1



- **False:** Even if your patient does not abuse their medication, accidental overdoses can happen and naloxone is an important safety precaution that helps keep them and their loved ones safe.
 - While your patient may not seem like they are at risk for an overdose, having a RIOSORD > 32 indicates that they may have a combination of smaller risk factors that puts them at a greater overall risk.

Question 2



True or False?

- If I inform patients that naloxone is available, this will **not** encourage them to abuse drugs.

Question 2



- **Truth:** Studies report that naloxone does not encourage drug use. In some cases, naloxone has been shown to decrease drug use. Naloxone blocks the effects of opiates and can produce unpleasant withdrawal symptoms.
 - Following a successful overdose reversal, a patient can access additional treatment options that he or she may not have considered previously.

Question 3

True or False?

- Naloxone is difficult to use.

Question 3



- **False:** Naloxone comes in several forms. We generally recommend the intranasal form (e.g., Narcan) which allows people to spray naloxone into the patient’s nostrils. Distribute the “Naloxone Administration” brochure to walk through the process with the patient.
- We recommend calling 911, administering the first dose, and then administering a second dose if the patient is not breathing 2-3 minutes after the first dose; or responds to the first dose but stops breathing again. Naloxone wears off after 30 to 60 minutes.

Question 4

True or False?

- My patients that are active duty service members will be flagged or placed on a “list” if they are co-prescribed naloxone.

Question 4



- **False:** The policy for administering naloxone applies to the entire MHS. MTF Commanders should ensure that their MTF implements this policy and service members should not encounter any issues for having a naloxone prescription.

Question 5



True or False?

- Clinical providers do not need to write a prescription for a patient to receive naloxone.

Question 5



- **True:** DHA-PI 6025.07 for “Naloxone Prescribing and Dispensing by Pharmacists in Military Treatment Facilities” authorizes pharmacists to dispense naloxone upon patient request.

Additional Resources



- The OEND page provides additional resources for patients, caregivers, prescribers, and pharmacists, such as:
 - Training Resources
 - CarePoint Instructions
 - The Quick Reference Guide

health.mil/opioidsafety

A screenshot of the Health.mil website. The page title is "Opioid Safety" under the "Conditions and Treatments" section. The website header includes the Health.mil logo, navigation links (Home, About the MHS, Topics, Training, Policies, Reference Center, News & Gallery, I am a...), and a search bar. The main content area is titled "Opioid Safety" and contains text about managing pain, a list of options (acetaminophen, exercise therapy, cognitive behavioral therapy), and information about common prescription opioids. A sidebar on the left lists various conditions and treatments, with "Opioid Safety" selected. A right sidebar contains links for "Prevent Opioid Misuse", "Rx Awareness", and "More CDC Information".

Key Takeaways



- Prescribing naloxone is standard best practice
- The Quick Reference Guide helps determine when and how to prescribe naloxone:
 - Assess if your patients are at risk for an overdose
 - Educate your patients (and their family member/support person when possible) about opioid safety and naloxone administration
 - Document elevated risk indications and naloxone discussion and prescription in your patient's electronic health record

Any patient that meets the following risk criteria should be prescribed naloxone:

- RIOSORD>32
- MEDD>50
- Opioid/Benzodiazepine combination use
- On long-term opioid therapy

Key Takeaways (Cont.)



- Talking to patients about naloxone:
 - Use key messages
 - Educate both patients and their caregivers
 - Be prepared to answer questions
- Share resources with other members of your healthcare team
- Work with your team to develop a day-to-day action plan for educating and co-prescribing naloxone to patients

For more information, visit health.mil/opioidsafety

References



- Defense Health Agency. (June 19, 2018). *Naloxone Prescribing and Dispensing by Pharmacists in Medical Treatment Facilities*. (DHA-PI 6025.07). Retrieved from <https://health.mil/Reference-Center/Policies/2018/06/19/DHA-PI-6025-07-Naloxone-in-the-MTFs>
- Department of Veterans Affairs (2017). VA/DoD clinical practice guideline for opioid therapy for chronic pain. Retrieved from <https://www.healthquality.va.gov/guidelines/Pain/cot/VADoDOTCPG022717.pdf>
- Murphy, L., Babaei-Rad, R., Buna, D., Isaac, P., Murphy, A., Ng, K., ... & Sproule, B. (2018). Guidance on opioid tapering in the context of chronic pain: Evidence, practical advice and frequently asked questions. *Canadian Pharmacists Journal/Revue des Pharmaciens du Canada*, 151(2), 114-120. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5843113/>

Questions?