

## References and Acknowledgements:

This clinical recommendation was developed on the basis of a thorough literature search by the Traumatic Brain Injury Center of Excellence (TBICoE) core working group, and was supported by the consensus of an expert working group and by the input of an end-user working group. A full literature search is available upon request. The TBICoE team wishes to acknowledge the contributions of the members of the expert and end-user working groups, listed below, and express our sincere gratitude. Many thanks.

### Expert Working Group:

Alicia Souvigner, DPT, CAPT, USN

Anne Mucha, DPT, MS, NCS

Anthony Kontos, PhD

Carrie Hoppes, DPT, PhD, LTC, USA

Christopher Giza, MD

F. J. Haran, PhD, LCDR, USN

John Leddy, MD

Karen McCulloch, DPT, PhD

Kimberly G. Harmon, MD

Michael McCrea, PhD

Noah Silverberg, PhD

Scott Pyne, MD, CAPT, USN

Steven Porter, PsyD, LCDR, USN

Wesley Cole, PhD

### End-User Working Group:

Adam Susmarski DO, CDR, MC, USN

Alejandro Navarro, PA-C, MAJ, USA-NG

Blaise Pascale, MD, MPH, MAJ, USA

Brian Colsant, MD

Brian Hwang, MD, SGT, USA

Daniel Hubbard, PA-C, MAJ USAF

Jessicah Ray, DHSc, MS, PA-C

Matthew Puderbaugh DO, Maj, USAF

Mellanie Medina, FNP-BC

Michael Collins, HM1, USN, NREMT

Michael Winkle, HMC, IDC, USN

Michelle Lea, PA-C, LCDR, USN

Michelle Lindsay, ANP-BC

Scott Pyne, MD, CAPT, USN

Stephanie Maxfield-Panker, DPT, PhD

Tyler Bates, DO, Capt, USAF

Zachary Wilson, HM1, IDC, USN

### TBICoE Core Working Group:

Amanda Gano, MS, PA-C

Donald Marion, MD, MSc

Gary McKinney, DHSc, CBIS

Joanne Gold, PharmD, BCGP

Keith Stuessi, MD

Stacey Harcum, MPH, MS, OTR/L

## References:

- Adam O, Mac Donald CL, Rivet D, et al. Clinical and imaging assessment of acute combat mild traumatic brain injury in Afghanistan. *Neurology*. 2015;85(3):219-227.
- Alsalaheen B, Stockdale K, Pechumer D, Broglio SP. Measurement Error in the Immediate Postconcussion Assessment and Cognitive Testing (ImPACT): Systematic Review. *J Head Trauma Rehabil*. 2016;31(4):242-251.
- ANAM. Automated Neurocognitive Assessment Metric. 2019; <https://armymedicine.health.mil/Regional-Health-Commands/Office-of-the-Surgeon-General/R2D/ANAM>, 2019.
- Asken BM, Bauer RM, Guskiewicz KM, et al. Immediate Removal From Activity After Sport-Related Concussion Is Associated With Shorter Clinical Recovery and Less Severe Symptoms in Collegiate Student-Athletes. *Am J Sports Med*. 2018;46(6):1465-1474.
- Asken BM, Clugston JR, Snyder AR, Bauer RM. Baseline Neurocognitive Performance and Clearance for Athletes to Return to Contact. *J Athl Train*. 2017;52(1):51-57.
- Asken BM, McCrea MA, Clugston JR, Snyder AR, Houck ZM, Bauer RM. "Playing Through It": Delayed Reporting and Removal From Athletic Activity After Concussion Predicts Prolonged Recovery. *J Athl Train*. 2016;51(4):329-335.
- Baillie JM, Remigio-Baker RA, Cole WR, et al. Use of the Progressive Return to Activity Guidelines May Expedite Symptom Resolution After Concussion for Active Duty Military. *Am J Sports Med*. 2019;47(14):3505-3513.
- Black AM, Sergio LE, Macpherson AK. The Epidemiology of Concussions: Number and Nature of Concussions and Time to Recovery Among Female and Male Canadian Varsity Athletes 2008 to 2011. *Clin J Sport Med*. 2017;27(1):52-56.
- Broglio SP, Macciocchi SN, Ferrara MS. Neurocognitive performance of concussed athletes when symptom free. *Journal of athletic training*. 2007;42(4):504-508.
- Brooks BL, Low TA, Daya H, Khan S, Mikrogianakis A, Barlow KM. Test or Rest? Computerized Cognitive Testing in the Emergency Department after Pediatric Mild Traumatic Brain Injury Does Not Delay Symptom Recovery. *J Neurotrauma*. 2016;33(23):2091-2096.
- Brown NJ, Mannix RC, O'Brien MJ, Gostine D, Collins MW, Meehan WP, 3rd. Effect of cognitive activity level on duration of post-concussion symptoms. *Pediatrics*. 2014;133(2):e299-304.
- Cole WR, Arrieux JP, Ivins BJ, Schwab KA, Qashu FM. A Comparison of Four Computerized Neurocognitive Assessment Tools to a Traditional Neuropsychological Test Battery in Service Members with and without Mild Traumatic Brain Injury. *Archives of clinical neuropsychology : the official journal of the National Academy of Neuropsychologists*. 2018;33(1):102-119.

Cole WR, Arrieux JP, Schwab K, Ivins BJ, Qashu FM, Lewis SC. Test-retest reliability of four computerized neurocognitive assessment tools in an active duty military population. *Archives of clinical neuropsychology: the official journal of the National Academy of Neuropsychologists*. 2013;28(7):732-742.

Cottle JE, Hall EE, Patel K, Barnes KP, Ketcham CJ. Concussion Baseline Testing: Preexisting Factors, Symptoms, and Neurocognitive Performance. *J Athl Train*. 2017;52(2):77-81.

Covassin T, Elbin RJ, 3rd, Stiller-Ostrowski JL, Kontos AP. Immediate post-concussion assessment and cognitive testing (ImPACT) practices of sports medicine professionals. *J Athl Train*. 2009;44(6):639-644.

Daniel JC, Olesniewicz MH, Reeves DL, et al. Repeated measures of cognitive processing efficiency in adolescent athletes: implications for monitoring recovery from concussion. *Neuropsychiatry, neuropsychology, and behavioral neurology*. 1999;12(3):167-169.

Darling SR, Leddy JJ, Baker JG, et al. Evaluation of the Zurich Guidelines and exercise testing for return to play in adolescents following concussion. *Clin J Sport Med*. 2014;24(2):128-133.

DiFazio M, Silverberg ND, Kirkwood MW, Bernier R, Iverson GL. Prolonged Activity Restriction After Concussion: Are We Worsening Outcomes? *Clin Pediatr (Phila)*. 2016;55(5):443-451.

Ebell MH, Siwek J, Weiss BD, et al. Strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature. *J Am Board Fam Pract*. 2004;17(1):59-67.

Echemendia RJ, Bruce JM, Meeuwisse W, Comper P, Aubry M, Hutchison M. Long-term reliability of ImPACT in professional ice hockey. *The Clinical Neuropsychologist*. 2016;30(2):328-337.

Elbin RJ, Sufrinko A, Schatz P, et al. Removal From Play After Concussion and Recovery Time. *Pediatrics*. 2016;138(3).

Gregory E, West TA, Cole WR, et al. Use of a multi-level mixed methods approach to study the effectiveness of a primary care progressive return to activity protocol after acute mild traumatic brain injury/concussion in the military. *Contemp Clin Trials*. 2017;52:95-100.

Group TMOc-mTBIW. VA/DoD Clinical Practice Guideline For The Management Of Concussion-Mild Traumatic Brain Injury. 2016;  
<https://www.healthquality.va.gov/guidelines/Rehab/mtbi/mTBICPGFullCPG50821816.pdf>, 2019.

Guskiewicz KM, Broglio SP. Acute sports-related traumatic brain injury and repetitive concussion. *Handbook of Clinical Neurology*. 2015;127:157-172.

Haran FJ, Schumacher P, Markwald R, Handy JD, Tsao JW. Relationships Between Sleepiness, Mood, and Neurocognitive Performance in Military Personnel. *Front Neurol*. 2019;10:674.

Harmon KG, Drezner JA, Gammons M, et al. American Medical Society for Sports Medicine position statement: concussion in sport. *Br J Sports Med*. 2013;47(1):15-26.

Huang M, Frantz J, Morales G, et al. Reduced Resting and Increased Elevation of Heart Rate Variability With Cognitive Task Performance in Concussed Athletes. *The Journal of Head Trauma Rehabilitation*. 2019;34(1):45-51.

Kelley AM, Raney BM, Estrada A, Grandizio CM. Evaluation of the Military Functional Assessment Program: Preliminary Assessment of the Construct Validity Using an Archived Database of Clinical Data. *The Journal of head trauma rehabilitation*. 2015;30(4):E11-20.

Kelley AM, Showers M. Evaluation of the Military Functional Assessment Program for Return-to-Duty Decision Making: A Longitudinal Study. *Mil Med*. 2019;184(Suppl 1):160-167.

Lawrence DW, Richards D, Comper P, Hutchison MG. Earlier time to aerobic exercise is associated with faster recovery following acute sport concussion. *PloS one*. 2018;13(4):e0196062.

Leddy JJ, Haider MN, Ellis MJ, et al. Early Subthreshold Aerobic Exercise for Sport-Related Concussion: A Randomized Clinical Trial. *JAMA Pediatr*. 2019;173(4):319-325.

Leddy JJ, Wilber CG, Willer BS. Active recovery from concussion. *Curr Opin Neurol*. 2018;31(6):681-686.

Leddy JJ, Willer B. Use of Graded Exercise Testing in Concussion and Return-to-Activity Management. *Current Sports Medicine Reports*. 2013;12(6):370-376.

Lee YM, Wu A, Zuckerman SL, et al. Obesity and neurocognitive recovery after sports-related concussion in athletes: a matched cohort study. *The Physician and Sports Medicine*. 2016;44(3):217-222.

Luethcke CA, Bryan CJ, Morrow CE, Isler WC. Comparison of concussive symptoms, cognitive performance, and psychological symptoms between acute blast-versus nonblast-induced mild traumatic brain injury. *Journal of the International Neuropsychological Society : JINS*. 2011;17(1):36-45.

Manderino LM, Gunstad J. Performance of the Immediate Post-Concussion Assessment and Cognitive Testing Protocol Validity Indices. *Archives of clinical neuropsychology : the official journal of the National Academy of Neuropsychologists*. 2018;33(5):596-605.

Mayers LB, Redick TS. Clinical utility of ImpACT assessment for postconcussion return-to-play counseling: psychometric issues. *Journal of clinical and experimental neuropsychology*. 2012;34(3):235-242.

McClincy MP, Lovell MR, Pardini J, Collins MW, Spore MK. Recovery from sports concussion in high school and collegiate athletes. *Brain Inj*. 2006;20(1):33-39.

McClure DJ, Zuckerman SL, Kutscher SJ, Gregory AJ, Solomon GS. Baseline neurocognitive testing in sports-related concussions: the importance of a prior night's sleep. *Am J Sports Med*. 2014;42(2):472-478.

McCrea M, Broglio S, McAllister T, et al. Return to play and risk of repeat concussion in collegiate football players: comparative analysis from the NCAA Concussion Study (1999-2001) and CARE Consortium (2014-2017). *Br J Sports Med*. 2019.

Meyer JE, Arnett PA. Changes in symptoms in concussed and non-concussed athletes following neuropsychological assessment. *Dev Neuropsychol*. 2015;40(1):24-28.

Norris JN, Carr W, Herzig T, Labrie DW, Sams R. ANAM4 TBI reaction time-based tests have prognostic utility for acute concussion. *Mil Med*. 2013;178(7):767-774.

Orr R, Bogg T, Fyffe A, Lam LT, Browne GJ. Graded Exercise Testing Predicts Recovery Trajectory of Concussion in Children and Adolescents. *Clin J Sport Med*. 2018.

Pawlukiewicz A, Yengo-Kahn AM, Solomon G. The Effect of Pretest Exercise on Baseline Computerized Neurocognitive Test Scores. *Orthop J Sports Med*. 2017;5(10):2325967117734496.

Popovich M, Almeida A, Freeman J, et al. Use of Supervised Exercise During Recovery Following Sports-Related Concussion. *Clin J Sport Med*. 2019.

Porter SJ, Johnson DE. Clinical Use of the Automated Neuropsychological Assessment Metrics TBI-Mil Expanded Battery in Evaluating Concussion Recovery: A Retrospective Study. *Mil Med*. 2020;185(9-10):e1722-e1727. doi:10.1093/milmed/usaa075.

Randolph C. Baseline neuropsychological testing in managing sport-related concussion: does it modify risk? *Current sports medicine reports*. 2011;10(1):21-26.

Remigio-Baker RA, Bailie JM, Gregory E, et al. Activity Level and Type During Post-acute Stages of Concussion May Play an Important Role in Improving Symptoms Among an Active Duty Military Population. *Front Neurol*. 2019;10:602.

Sandel NK, Lovell MR, Kegel NE, Collins MW, Kontos AP. The relationship of symptoms and neurocognitive performance to perceived recovery from sports-related concussion among adolescent athletes. *Applied neuropsychology Child*. 2013;2(1):64-69.

Saviola D, De Tanti A, Conforti J, et al. Safe return to driving following severe acquired brain injury: role of a short neuropsychological assessment. *European journal of physical and rehabilitation medicine*. 2018;54(5):717-723.

Sours C, George EO, Zhuo J, Roys S, Gullapalli RP. Hyper-connectivity of the thalamus during early stages following mild traumatic brain injury. *Brain Imaging Behav*. 2015;9(3):550-563.

Stone S, Lee B, Garrison JC, Blueitt D, Creed K. Sex Differences in Time to Return-to-Play Progression After Sport-Related Concussion. *Sports Health*. 2017;9(1):41-44.

Thomas DG, Apps JN, Hoffmann RG, McCrea M, Hammeke T. Benefits of strict rest after acute concussion: a randomized controlled trial. *Pediatrics*. 2015;135(2):213-223.

Tsushima WT, Yamamoto MH, Ahn HJ, Siu AM, Choi SY, Murata NM. Invalid Baseline Testing with ImPACT: Does Sandbagging Occur with High School Athletes? *Applied neuropsychology Child*. 2019:1-10.

Wiebe DJ, Nance ML, Houseknecht E, et al. Ecologic Momentary Assessment to Accomplish Real-Time Capture of Symptom Progression and the Physical and Cognitive Activities of Patients Daily Following Concussion. *JAMA Pediatr*. 2016;170(11):1108-1110.

#### FAQ References:

1. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc.* 1982;14(5):377-381.
2. Lucas S. Headache management in concussion and mild traumatic brain injury. *PM&R.* Oct 2011;3(10 Suppl 2):S406-412.
3. Iverson GL, Gardner AJ, Terry DP, et al. Predictors of clinical recovery from concussion: a systematic review. *Br J Sports Med.* Jun 2017;51(12):941-948.
4. Collins MW, Kontos AP, Reynolds E, Murawski CD, Fu FH. A comprehensive, targeted approach to the clinical care of athletes following sport-related concussion. *Knee Surg Sports Traumatol Arthrosc.* Feb 2014;22(2):235-246.
5. Yue JK, Cnossen MC, Winkler EA, et al. Pre-injury Comorbidities Are Associated With Functional Impairment and Post-concussive Symptoms at 3- and 6-Months After Mild Traumatic Brain Injury: A TRACK-TBI Study. *Front Neurol.* 2019;10(343):343.
6. Scott BR, Uomoto JM, Barry ES. Impact of pre-existing migraine and other co-morbid or co-occurring conditions on presentation and clinical course following deployment-related concussion. *Headache.* 2020;0:1-16.
7. Iljazi A, Ashina H, Al-Khazali HM, et al. Post-Traumatic Stress Disorder After Traumatic Brain Injury-A Systematic Review and Meta-Analysis. *Neurol Sci.* May 15 2020.
8. DVBC. Research Review on Mild Traumatic Brain Injury and Posttraumatic Stress Disorder. 2018.