


Alcohol Misuse: Facts about Risky Drinking

How Much is Too Much?

It is easy to drink a lot without realizing it. A standard drink contains about 0.6 fluid ounces of pure alcohol, but many drinks, like a large 22-ounce beer or a mixed cocktail, contain more than one standard drink.

The image below highlights examples of what a standard drink looks like for different beverage types.



Risky Drinking

Risk levels for drinking are determined by the amount you drink during the week and on any single occasion. Use the chart below to determine your level of drinking.

Level of Drinking	Number of Drinks	
	Male 65 and Below	Male 65 and Above AND Female 65 and Below
Low risk	No more than 4 per day or 14 per week	No more than 3 per day or 7 per week
Heavy	More than 14 per week	More than 7 per week
Binge	5 or more per single occasion	4 or more per single occasion

Preventing Risky Drinking

Monitoring your alcohol use can help you prevent risky drinking:

- Set a daily and weekly drinking limit
- Pace your drinking
- Record how much you drink each day
- Avoid situations and people or places that cause you to drink
- Ask a friend who does not drink to help you stay within your limit

Indicators of Potential Alcohol Use Disorder

The box below lists behaviors associated with risky and problematic drinking. If you respond "yes" to any of the indicators in the past year, read the section on Where to Seek Help.

- Drink more, or longer, than you intend
- Try to cut down or stop drinking, but are not able to
- Have cravings or strong urges to have a drink
- Have to drink more than you once did to get the effect you want
- Continue to drink even though it makes you feel depressed or anxious or adds to another health problem
- Spend a lot of time drinking, getting alcohol, or recovering from drinking
- Find that drinking often interferes with daily activities, family, friends and/or work
- Cessation of or reduction in alcohol use that has heavy or long results with withdrawal symptoms. Such symptoms can include sweating or high pulse rate, hand tremor, insomnia, nausea or vomiting, hallucinations, agitation, anxiety, or seizure.

Where to Seek Help

If you think that you, or someone you know, may be engaging in risky drinking, seek help from medical providers, leaders, or battle buddies.

Below are some additional resources:

- Contact your primary care manager for referrals and medical assistance
- Military Health System – information about addiction, substances, treatments and getting help health.mil
- TRICARE Alcohol Awareness – information and support resources on excessive drinking in the military tricare.mil/HealthWellness/Alcohol
- U.S. Department of Veterans Affairs – For help making medical appointments and with prescription refills myhealth.va.gov
- Military OneSource – online, telephone, and face-to-face non-medical counseling militaryonesource.mil | (800) 342-9647
- Psychological Health Resource Center – a 24/7 anonymous information call center www.health.mil/PHRC | (866) 966-1020
- Veterans Crisis Line – a 24/7 anonymous crisis hot line <http://www.veteranscrisisline.net> | (800) 273-8255 and press "1"

Impact of Risky Drinking

Excessive drinking places your safety and health at risk. See below for some of the effects of risky drinking.



- Depression
- Aggressive behavior
- Anxiety
- Alcohol dependence
- Insomnia
- Memory loss



- Stroke
- Heart failure
- Hypertension
- Premature aging



- Frequent colds
- Reduced resistance to infection
- Increased risk of pneumonia



- Anemia
- Vitamin deficiency
- Blood clotting
- Bleeding



- Cancer of the throat and mouth
- Breast cancer
- Inflammation of the pancreas



- Stomach inflammation
- Diarrhea
- Malnutrition



- Risk of fetal alcohol spectrum disorders, which include physical, behavioral and learning disabilities



- Painful nerves
- Numb, tingling toes
- Impaired sensations leading to falls



- Type II Diabetes
- Liver damage



- Motor vehicle crashes
- Failure to fulfill obligations at work, school and home



- Men: Erectile dysfunction
- Women: Unintended pregnancy
- Sexually transmitted diseases



- Injury
- Legal problems
- Violence
- Impacts service member's readiness
- Violent crime

References

National Center on Birth Defects and Developmental Disabilities. (2014). *Planning and Implementing Screening and Brief Intervention for Risky Alcohol Use: A Step-by-Step Guide for Primary Care Practices*. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. <https://www.cdc.gov/ncbddd/fasd/documents/alcoholbsiimplementationguide.pdf>

Meadows, S. O., Engel, C. C., Collins, R. L., Beckman, R. L., Cefalu, M., Hawes-Dawson, J., Doyle, M., Kress, A. M., Sontag-Padilla, L., Ramchand, R., & Williams, K. M. (2018). 2015 Department of Defense Health Related Behaviors Survey (HRBS). *Rand health quarterly*, 8(2), 5. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6183770/>

National Institute on Alcohol Abuse and Alcoholism (NIAAA). (2010). *Rethinking Drinking: Alcohol and Your Health* (NIH Publication No. 13-3770). U.S. Department of Health and Human Services, National Institutes of Health. http://pubs.niaaa.nih.gov/publications/rethinkingdrinking/rethinking_drinking.pdf

National Institute on Alcohol Abuse and Alcoholism (NIAAA). (n.d.). *Rethinking Drinking: How much is too much?*. U.S. Department of Health and Human Services, National Institutes of Health. <https://www.rethinkingdrinking.niaaa.nih.gov/How-much-is-too-much/>

