Facility	
Position Title	Trainee Name
Assessment Start Date	Assessment Completion Date

This competency form is to be used as an addendum to initial/annual competency assessment checklist. This competency highlights the special rules for preparation and administration of JE-VC (Ixiaro) to pediatric patients.

	Self- Assessment* (Preceptor Date & Initials)		totion	Validation of Competency			
Required Competency or Skill			Evaluation Method+	Date	Initials	Comments	
Vaccine Administration Procedures	CRITICAL THINKING: Follows manufacturer guidelines. Administers the right vaccin to the right patient at the right time. Understands that the deviation from the recomm may reduce vaccine effectiveness or increase the risk of local reactions. ** Aspiration toxoids (i.e., pulling back on the syringe plunger after needle insertion, before injection large blood vessels exists at the recommended injection sites.					ine, right dose, a mended route of a on before injectio tion) is not require	nd right route, administration n of vaccines or ed because no
A. Selects appropriate JE-VC product based on age and screening of patient							
B. Verbalizes understanding of the standing order and package insert for the administration of JE-VC vaccine to adult and pediatric patients							
C. Gathers required supplies for administering JE-VC vaccine (i.e. gauze, alcohol pads, bandages, sharps container, etc.)							
D. Washes hands or uses hand sanitizer between patients							
E. Prepares JE-VC vaccine for administration							
1. Removes properly stored JE-VC vaccine product from storage unit							
a. Ensures vaccine was stored in refrigerator at (2-8°C)							
b. Inspects syringe for damage or contamination							
c. Checks vaccine(s) expiration date(s)							
2. Checks to make sure proper product is picked based on age							

* Self Assessment:	1=Experienced	2=Needs Practice/Ass	istance	3=Never Done	N/A=Not Applicable	
+ Evaluation / Validation	n Method:	T=Written Test		D=Demonstration/Observation	V=Verbal	I=Interactive Class
DHA-IHD (9 Aug 2024)			(877) GET-VACC (DSN 76	1-4245)	W	ww.health.mil/vaccines

		Salf		Validation of Competency					
		Required Competency or Skill	Assessment*	(Preceptor Date & Initials)		Evaluation Method+	Date	Initials	Comments
	a.	2 – 35 months use syringe with red marking on barrel							
	b.	36 months and older, use syringe with or without red marking on barrel							
3.	Pre adı	epares prefilled syringe for a 0.25ml dose to ninister JE vaccine to children 2-35 months of age							
	a.	Shake 0.5ml prefilled syringe to thoroughly mix contents							
	b.	Remove tip cap and attach appropriate size needle							
	C.	Holds the syringe in upright position and uncaps the needle							
	d.	Push the plunger stopper up to the edge of the red line on the syringe barrel, indicated by a red arrow (see figure below)*, and discards expelled volume into a medical waste container. Note: If plunger stopper is pushed beyond redline, do not administer the vaccine. Repeat the procedure using a new prefilled syringe							
	e.	Locks the needle safety shield and removes the needle							
	f.	Attaches a new sterile needle prior to injection of the remaining dose							
	g.	Ensures any manufacturer prefilled syringe with tip cap removed and needle attached is discard- ed at the end of duty day if not administered							

* Self Assessment:	1=Experienced	2=Needs Practice/Assistance	3	3=Never Done	N/A=Not Applica	ble
+ Evaluation / Validatio	n Methodologies:	T=Tests	C	D=Demonstration/Observation	V=Verbal	I=Interactive Class
DHA-IHD (9 Aug 2024)		(877) G	ET-VACC (DSN 761-	-4245)		www.health.mil/vaccines

		Calf	Self- Orientation ssment* (Preceptor Date & Initials)		Validation of Competency				
	Required Competency or Skill	Assessment*			Evaluation Method+	Date	Initials	Comments	
4.	Prepares prefilled syringe for a 0.5ml dose to administer JE vaccine to individuals 36 months and older								
	 Shake the prefilled syringe containing 0.5ml to thoroughly mix contents 								
	 Remove the syringe tip cap and attach appropriate size needle 								
5.	Maintains aseptic technique throughout vaccine preparation process								
F. A g in	dministers JE-VC vaccine per ACIP/manufacturer uidelines ensuring proper route, dosage, timing, and dications/ contraindications								
1.	Selects 22-25g needle and appropriate length based on administration route and body size								
2.	Selects appropriate dose based on age								
	a. 2-35 months = 0.25 ml								
	b. 36 months and older = 0.5 ml								
3.	Selects appropriate anatomical site based on age								
	 Infants and toddlers (lacking adequate deltoid mass); anterolateral aspect of thigh 								
	b. Toddler/Children/Teens/Adults: the deltoid muscle is recommended for routine intramuscular vaccinations; demonstrates 3 fingers down from acromion process to select proper area								
4.	Cleans injection site with alcohol								
5.	Inserts the needle fully into the muscle at a 90o angle (per ACIP aspiration is not required)								
6.	Injects vaccine using steady pressure then withdraws needle at angle of insertion								
7.	Applies light pressure with gauze to injection site for several seconds								
* Self A	ssessment: 1=Experienced 2=Needs Pra	actice/Assistance		3=Never Done N/A=Not Applicable		Applicable			
+ Evalu	Evaluation / Validation Methodologies: T=Tests D=Demonstration/Observation V=Verbal I=Interactive Class						eractive Class		
DHA-IH	DHA-IHD (9 Aug 2024) (877) GET-VACC (DSN 761-4245) www.health.mil/vacc					alth.mil/vaccines			

Preceptor's Initials	Printed Name	Signature

I understand the topics listed, I will be allowed to perform only those within scope of practice, and only after I have successfully my demonstrated competency.

Trainee Signature: _____

Date: _____



This document is not valid without a digital signature.