

HEALTH AFFAIRS

AUG 2 6 2010

The Honorable Carl Levin Chairman, Committee on Armed Services United States Senate Washington, DC 20510

Dear Mr. Chairman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charles L.K.

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable John McCain Ranking Member



HEALTH AFFAIRS

AUG 2 6 2010

The Honorable Ike Skelton Chairman, Committee on Armed Services U.S. House of Representatives Washington, DC 20515

Dear Mr. Chairman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charles L. R. -

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable Howard P. "Buck" McKeon Ranking Member



AUG 2 6 2010

HEALTH AFFAIRS

The Honorable Susan Davis Chairwoman, Subcommittee on Military Personnel Committee on Armed Services U.S. House of Representatives Washington, DC 20515

Dear Madam Chairwoman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charle L. R. -

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable Joe Wilson Ranking Member



HEALTH AFFAIRS

AUG 2 6 2010

The Honorable Daniel K. Inouye Chairman, Subcommittee on Defense Committee on Appropriations United States Senate Washington, DC 20510

Dear Mr. Chairman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charla L. R. -

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable Thad Cochran Ranking Member



HEALTH AFFAIRS

AUG 2 6 2010

The Honorable David R. Obey Chairman, Committee on Appropriations U.S. House of Representatives Washington, DC 20515

Dear Mr. Chairman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charles L. Kin

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable Jerry Lewis Ranking Member



AUG 2 6 2010

HEALTH AFFAIRS

The Honorable Norm Dicks Chairman, Subcommittee on Defense Committee on Appropriations U.S. House of Representatives Washington, DC 20515

Dear Mr. Chairman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charle C.R.

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable C. W. Bill Young Ranking Member



HEALTH AFFAIRS

AUG 2 6 2010

The Honorable James H. Webb Chairman, Subcommittee on Personnel Committee on Armed Services United States Senate Washington, DC 20510

Dear Mr. Chairman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charle L. R.

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable Lindsey O. Graham Ranking Member



HEALTH AFFAIRS

AUG 2 6 2010

The Honorable Daniel K. Inouye Chairman, Committee on Appropriations United States Senate Washington, DC 20510

Dear Mr. Chairman:

This report responds to House Report 110-652 of the National Defense Authorization Act for Fiscal Year 2009. The House Report requests the Secretary of Defense to study the possibility of providing a referral for a second opinion to potentially suicidal Service members in a combat theater.

Thank you for your continued support of the Military Health System.

Sincerely,

Charle L.R. -

Charles L. Rice, M.D. President, Uniformed Services University of the Health Sciences Performing the Duties of the Assistant Secretary of Defense (Health Affairs)

Enclosure: As stated

cc: The Honorable Thad Cochran Ranking Member



SUICIDE PREVENTION IN THE ARMED FORCES STUDY ON REFERRAL FOR SECOND OPINION

REPORT TO CONGRESS

HOUSE REPORT 110-652 OF THE NATIONAL DEFENSE AUTHORIZATION ACT FOR FISCAL YEAR 2009

FEBRUARY 2010

TABLE OF CONTENTS

INTRODUCTION
SECOND OPINION AND CONSULTATIONS: DEFINITION OF TERMS
DISCUSSION OF CONSULTATION VERSUS SECOND OPINION IN SUICIDE RISK ASSESSMENT4
DEPARTMENT OF DEFENSE POLICIES THAT GUIDE MENTAL HEALTH CARE
CONCLUSION
OVERVIEW OF THE LITERATURE

Introduction

This report responds to House Report 110-652, page 384, of the National Defense Authorization Act for Fiscal Year 2009 that directs the Secretary of Defense to study the possibility of providing a referral for second opinion to potentially suicidal Service members in the combat theater. In April 2009, OASD (HA) established the Suicide Prevention and Risk Reduction Committee (SPARRC). The SPARRC is chaired by the Defense Centers of Excellence (DCoE) and has representatives from the Army, Navy, Air Force, Marine Corps, Coast Guard, the National Guard Bureau Director of Psychological Health and Reserve Affairs, the Armed Forces Medical Examiner, the Armed Forces Chaplains Board, the DCoE Telehealth and Technology center, as well as, Veterans Affairs, Centers for Disease Control and Substance Abuse and Mental Health Administration. To provide an informed response, the SPARRC formed a working group to examine the topic. The following organizations participated in the working group:

Organization	
Air Force	
Defense Centers of Excellence for Psychological Health	and
Traumatic Brain Injury	
Uniformed Services University of Health Sciences	
Substance Abuse and Mental Health Services Administr	ation

The main focus of this report is the possibility of providing a second opinion for potentially suicidal Service members. The reviews and evaluations have resulted in ongoing discussions and changes to suicide prevention efforts. The working group believes professional consultation rather than second opinion is the preferred and more appropriate approach to clinical suicide risk assessment and management. The working group agreed that provider training and judgment should dictate when consultation is necessary. Consultation is already a common practice among providers. At this time, DoD has decided not to develop a policy to mandate consultation for suicidal patients in theater. However, the DoD Instruction on Combat Operational Stress Control will be modified to include language encouraging providers to consult their colleagues in evaluating patients who present a threat to self/others in the deployed environment. We will use this report to help guide and inform our current programs and policies.

Second Opinion and Consultations: Definition of Terms

Different definitions of consultation and second opinion may apply in various settings. For the purposes of this report, the working group adopted the following definitions (referenced throughout the report) to pertain specifically to communications among mental health providers. The primary objective of clinical communication is to make the most appropriate clinical care decisions for our Service members recognized at risk for suicide behavior.

A **Consultation** within the military mental health field is a provider request for input from another knowledgeable health professional, based on the clinical provider's clinical description of a case. In the mental health field, consultation usually occurs when the primary mental health provider confers with a professional colleague about the patient. The clinical consultant may or may not interact directly with the patient in the course of the professional consultation. The provider seeking consultation is usually fully responsible for the handling of the case in terms of diagnosis and treatment. In the military this can depend on the relationship and qualifications of the consultant. Requests for consultations may occur when a provider seeks to enhance the assessment and treatment he or she is providing to a patient. Consultations are often sought in both medical settings for a variety of reasons to ensure optimal assessment and treatment for a given individual. In some cases, the consultation may result in a request for a second opinion.

In the field of mental health, a **second opinion** differs from consultation and is generally a patient-initiated, full and independent evaluation by a second provider who directly interacts with the patient. Responsibility for clinical care determined by the patient's choice of the professional opinion. A request for a second opinion, for instance, may occur if an individual is given a psychiatric diagnosis and subsequently, chooses to request an independent evaluation by a second provider for confirmation, or refutation of the initial diagnosis. In this vein, a second opinion may be sought after diagnosis to confirm a recommended course of treatment or generate possible alternative treatments. Thus, second opinions are often sought in both medical settings to confirm or clarify a diagnosis and/or possible courses of treatment.

Discussion of Consultation versus Second Opinion in Suicide Risk Assessment

Based on collaboration with subject matter experts and several group discussions involving both civilian and military mental health providers, the working group recognizes that, in general, professional consultation is the preferred, more appropriate, and the less intrusive approach to clinical suicide risk assessment and management. Seeking a timely or contemporaneous clinical consultation is widely accepted in clinical suicidology practice for providers who work with suicidal individuals, and will ultimately help to ensure proper and appropriate medical and psychiatric decisions.

It is important to note the benefits of professional consultation, especially in the assessment, management, and treatment of suicidal behaviors. First, consultation presents the opportunity to further examine the details of a given case with another colleague. This helps to minimize further trauma to the Service member who would have to re-live the experience in the case of seeking a second opinion. Second, consultation provides reassurance to the suicidal individual and his/her family members and commands that the best possible care is being provided in order to prevent any potential

harm. It provides the Service member and his or her family a more informed medical opinion on issues such as return to duty, evaluation, hospitalization, and treatment options. There is no guarantee providers can assess the risk of suicide with 100 percent certainty, but nonetheless the diagnosis is their responsibility. Consultations enable providers to share experiences and gain an improved understanding of suicide risk to help them more accurately assess and mitigate risk. Third, a consultation has fewer logistical difficulties because it can be obtained in an easier manner, particularly in very rural or rugged terrains (i.e., telephone, chat or video teleconference). Since a consultation does not require the consulting provider to have an interaction with the patient, there are fewer scheduling constraints that would delay the provision of care. In comparison, a referral for a second opinion in theater has various logistical challenges that make it a more complicated option than consultation.

Given the factors outlined above, the working committee decided it was more logical to focus on determining the feasibility of consultation rather than second opinion in theater. However, in circumstances when the primary treatment provider does not have the expertise to adequately diagnose or treat the Service member's illness, a second opinion may be more appropriate than a consultation.

Conclusion

Based on the review, DoD recognizes that while a consultation is often more feasible, there may be situations where a second opinion is called for. DoD believes that second opinion should still remain an option for patients to initiate as they feel necessary, however the working group at this time believes consultation is the recommended approach to clinical suicide risk in theater assessment but a provider's training and judgment should dictate when consultation is appropriate and necessary. Consultation is already a common practice among providers. At this time, DoD has decided not to develop a policy to mandate consultation or second opinion for suicidal patients in theater. However, the DoD Instruction on Combat Operational Stress Control will be modified to include language encouraging providers to work with their colleagues in evaluating patients who present a threat to self/others in the deployed environment.

Overview of the Literature

The working group collected recent literature including information about consultation and second opinion in order to identify current practices, as well as possible models or programs that could be used in theater to reduce suicide. These references are listed below.

Air Force Medical Operations Agency, Population Health Support Division. (Unpublished manuscript). Air Force Guide for Managing Suicidal Behavior: Strategies, Resources, and Tools. 8320 Laser Rd., Brooks AFB, Texas.

Carroll, D. W. (2008). Perspectives in psychiatric consultation liaison nursing. *Perspectives in Psychiatric Care, 44*, 211-215.

Jobes, D. A. (2006). *Managing Suicidal Risk: A Collaborative Approach*. New York: The Guilford Press.

Jobes, D.A. & Berman, A.L. (1993). Suicide and malpractice liability: Assessing and revising policies, procedures, and practice in outpatient settings. *Professional Psychology: Research and Practice*, 24, 91-99.

Magne-Ingvar, U & Ojehagen, A. (1999). Significant others of suicide attempters: Their views at the time of the acute psychiatric consultation. *Social Psychiatry and Psychiatric Epidemiology*, 34, 73-79.

Matthieu, M. M., Cross, W., Batres, A. R., Flora, C. M., & Knox, K. L. (2008). Evaluation of gatekeeper training for suicide prevention in Veterans. *Archives of Suicide Research*, 12, 148-154.

Macdonald, E. A. (1997). Personal view second opinions in oncology: Nuisance or opportunity? *Clinical Oncology*, 9, 418-420.

McNiel, D. E., Fordwood, S. R., Weaver, C.M., Chamberlain, J. R., Hall, S. E., & Binder, R. L. (2008). Effects of training on suicide risk assessment. *Psychiatric Service*, 59, 1462-1465.

Oordt, M. S., Jobes, D. A., Fonseca, V. P., & Schmidt, S. M. (2009). Training mental health professionals to assess and manage suicidal behavior: Can provider confidence and practice behaviors be altered? *Suicide and Life-Threatening Behavior*, *39*, 21-32.

Oordt M. S., Jobes, D. A., Rudd, M. D., Fonseca, V. P., Runyan, C. N., Stea, J. B., Campise, R. L., & Talcott, G. W. (2005). Development of a clinical guide to enhance care for suicidal patients. *Professional Psychology: Research and Practice in the Public Domain, 36*, 208–218. Payne, S.E., Hill, J. V., & Johnson, D. E. (2008). The use of unit watch or command interest profile in the management of suicide and homicide risk: Rationale and guidelines for the military mental health professional. *Military Medicine*, *173*, 25-35.

Peterson, A. L., Baker, M. T., & McCarthy, K. R. (2008). Combat stress casualties in Iraq. Part 1: Behavioral health consultation at an expeditionary medical group. *Perspectives in Psychiatric Care, 44*, 146-158.

Peterson, A. L., Baker, M. T., & McCarthy, K. R. (2008). Combat stress casualties in Iraq. Part 2: Psychiatric screening prior to aeromedical evacuation. *Perspectives in Psychiatric Care*, 44, 159-168.

Rudd, M. D., Cukrowicz, K. C., & Bryan, C. J. (2008). Core competencies in suicide risk assessment and management: Implications for supervision. *Training and Education in Professional Psychology*, 2, 219-228.

Simpson, S. & Stacy, M. (2004). Avoiding the malpractice snare: Documenting suicide risk assessment. *Journal of Psychiatric Practice*, 10, 185-189.

Young, J. J., Ursano, R. J., Bally, R. E., & McNeil, D. S. (1989). Consultation to a clinic following suicide. *American Journal of Orthopsychiatry*, 59, 473-476.