

John F. Brundage, MD, MPH (COL, USA, Ret); Stephen B. Taubman, PhD; Devin J. Hunt, MS; Leslie L. Clark, PhD, MS

TABLE 4 a. Attributable incidence rates and attributable fractions of traumatic brain injury (TBI) diagnoses within 24 months after return from Iraq/Afghanistan, by determinants of likelihood of diagnosis, active component, U.S. Armed Forces, by follow-up period, April 2003–December 2013

a. TBI		Determinants of likelihood of diagnosis	2003–2005	2005–2007	2007–2009	2009–2011	2011–2012	2012–2013	
Overall	Active war service								
	Attributable incidence, % of returnees		0.71%	2.34%	4.96%	4.69%	4.63%	3.86%	
	Attributable fraction, % of diagnoses		13.1	32.6	44.8	39.0	36.1	34.8	
	Enhanced ascertainment								
	Attributable incidence, % of returnees		0.47%	0.60%	1.86%	3.10%	3.97%	2.99%	
	Attributable fraction, % of diagnoses		8.7	8.4	16.8	25.8	30.9	27.0	
Background	Attributable incidence, % of returnees		4.24%	4.24%	4.24%	4.24%	4.24%	4.24%	
	Attributable fraction, % of diagnoses		78.2	59.1	38.3	35.2	33.0	38.2	
	Service	Active war service							
		Attributable incidence, % of returnees		0.81%	3.28%	8.06%	5.32%	4.93%	4.61%
		Attributable fraction, % of diagnoses		13.6	37.9	50.4	33.9	30.0	31.6
		Enhanced ascertainment							
Attributable incidence, % of returnees		0.50%	0.72%	3.26%	5.73%	6.82%	5.31%		
Attributable fraction, % of diagnoses		8.4	8.3	20.4	36.5	41.6	36.4		
Army	Background								
	Attributable incidence, % of returnees		4.66%	4.66%	4.66%	4.66%	4.66%	4.66%	
	Attributable fraction, % of diagnoses		78.1	53.8	29.2	29.7	28.4	32.0	
	Navy	Active war service							
		Attributable incidence, % of returnees		0.56%	1.98%	1.90%	3.64%	3.06%	3.89%
		Attributable fraction, % of diagnoses		12.9	38.7	33.2	52.1	42.5	49.7
Enhanced ascertainment									
Attributable incidence, % of returnees		0.79%	0.14%	0.84%	0.35%	1.15%	0.94%		
Attributable fraction, % of diagnoses		18.2	2.7	14.7	5.0	16.0	12.0		
Air Force	Background								
	Attributable incidence, % of returnees		2.99%	2.99%	2.99%	2.99%	2.99%	2.99%	
	Attributable fraction, % of diagnoses		68.9	58.5	52.2	42.8	41.5	38.2	
	Marine Corps	Active war service							
		Attributable incidence, % of returnees		0.36%	0.48%	1.07%	0.45%	1.29%	0.86%
		Attributable fraction, % of diagnoses		7.5	9.6	19.1	7.7	20.5	15.0
Enhanced ascertainment									
Attributable incidence, % of returnees		0.72%	0.79%	0.81%	1.65%	1.27%	1.17%		
Attributable fraction, % of diagnoses		15.0	15.8	14.5	28.4	20.2	20.3		
Background	Attributable incidence, % of returnees		3.72%	3.72%	3.72%	3.72%	3.72%	3.72%	
	Attributable fraction, % of diagnoses		77.5	74.5	66.4	63.9	59.2	64.7	
	Sex	Active war service							
		Attributable incidence, % of returnees		0.93%	2.06%	2.48%	4.07%	5.18%	2.40%
		Attributable fraction, % of diagnoses		17.1	28.2	28.7	35.6	41.9	25.7
		Enhanced ascertainment							
Attributable incidence, % of returnees		0.01%	0.74%	1.65%	2.84%	2.67%	2.43%		
Attributable fraction, % of diagnoses		0.2	10.1	19.1	24.9	21.6	26.0		
Male	Background								
	Attributable incidence, % of returnees		4.51%	4.51%	4.51%	4.51%	4.51%	4.51%	
	Attributable fraction, % of diagnoses		82.8	61.7	52.2	39.5	36.5	48.3	
	Female	Active war service							
		Attributable incidence, % of returnees		0.66%	2.47%	5.27%	5.09%	5.05%	4.15%
		Attributable fraction, % of diagnoses		11.7	32.5	45.3	40.3	37.6	35.9
Enhanced ascertainment									
Attributable incidence, % of returnees		0.53%	0.67%	1.92%	3.09%	3.93%	2.96%		
Attributable fraction, % of diagnoses		9.4	8.8	16.5	24.5	29.3	25.6		
Background	Attributable incidence, % of returnees		4.45%	4.45%	4.45%	4.45%	4.45%	4.45%	
	Attributable fraction, % of diagnoses		78.9	58.6	38.2	35.2	33.1	38.5	
	Age group	Active war service							
		Attributable incidence, % of returnees		0.14%	0.40%	1.63%	1.10%	0.77%	0.91%
		Attributable fraction, % of diagnoses		4.0	10.0	25.0	14.7	9.5	12.6
		Enhanced ascertainment							
Attributable incidence, % of returnees		0.38%	0.58%	1.87%	3.37%	4.36%	3.32%		
Attributable fraction, % of diagnoses		10.8	14.5	28.7	45.1	53.6	45.9		
<25 years	Background								
	Attributable incidence, % of returnees		3.01%	3.01%	3.01%	3.01%	3.01%	3.01%	
	Attributable fraction, % of diagnoses		85.3	75.4	46.2	40.2	37.0	41.6	
	Background	Active war service							
		Attributable incidence, % of returnees		0.79%	2.79%	5.36%	5.58%	5.79%	4.98%
		Attributable fraction, % of diagnoses		14.9	37.0	47.0	44.1	43.7	44.9
Enhanced ascertainment									
Attributable incidence, % of returnees		0.65%	0.90%	2.19%	3.21%	3.60%	2.26%		
Attributable fraction, % of diagnoses		12.3	11.9	19.2	25.4	27.2	20.4		
Background	Attributable incidence, % of returnees		3.85%	3.85%	3.85%	3.85%	3.85%	3.85%	
	Attributable fraction, % of diagnoses		72.8	51.1	33.8	30.5	29.1	34.7	

Table 4a (cont.)

Determinants of likelihood of diagnosis		2003–2005	2005–2007	2007–2009	2009–2011	2011–2012	2012–2013
25–35 years	Active war service						
	Attributable incidence, % of returnees	0.68%	2.20%	5.26%	4.40%	4.16%	3.20%
	Attributable fraction, % of diagnoses	12.1	31.0	47.7	37.8	33.3	29.6
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.50%	0.47%	1.34%	2.81%	3.93%	3.18%
	Attributable fraction, % of diagnoses	8.9	6.6	12.2	24.2	31.4	29.4
>35 years	Background						
	Attributable incidence, % of returnees	4.42%	4.42%	4.42%	4.42%	4.42%	4.42%
	Attributable fraction, % of diagnoses	78.9	62.3	40.1	38.0	35.3	40.9
	Active war service						
	Attributable incidence, % of returnees	0.50%	1.52%	3.11%	3.23%	2.86%	2.67%
	Attributable fraction, % of diagnoses	9.5	23.3	30.6	27.9	22.4	22.3
Race/ethnicity	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.12%	0.36%	2.40%	3.72%	5.25%	4.66%
	Attributable fraction, % of diagnoses	2.3	5.5	23.6	32.1	41.2	38.9
	Background						
	Attributable incidence, % of returnees	4.64%	4.64%	4.64%	4.64%	4.64%	4.64%
	Attributable fraction, % of diagnoses	88.2	71.2	45.7	40.0	36.4	38.8
White, non-Hispanic	Active war service						
	Attributable incidence, % of returnees	0.60%	2.40%	5.10%	4.74%	4.91%	4.06%
	Attributable fraction, % of diagnoses	10.6	31.3	44.1	37.7	36.9	35.8
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.56%	0.75%	1.94%	3.30%	3.87%	2.76%
	Attributable fraction, % of diagnoses	9.9	9.8	16.8	26.3	29.1	24.3
Black, non-Hispanic	Background						
	Attributable incidence, % of returnees	4.52%	4.52%	4.52%	4.52%	4.52%	4.52%
	Attributable fraction, % of diagnoses	79.6	58.9	39.1	36.0	34.0	39.9
	Active war service						
	Attributable incidence, % of returnees	0.72%	1.34%	3.50%	3.38%	3.14%	3.08%
	Attributable fraction, % of diagnoses	15.1	23.2	37.3	31.6	27.7	29.6
Hispanic	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.25%	0.63%	2.08%	3.50%	4.38%	3.53%
	Attributable fraction, % of diagnoses	5.2	10.9	22.2	32.8	38.7	33.9
	Background						
	Attributable incidence, % of returnees	3.80%	3.80%	3.80%	3.80%	3.80%	3.80%
	Attributable fraction, % of diagnoses	79.7	65.9	40.5	35.6	33.6	36.5
Other/unspecified	Active war service						
	Attributable incidence, % of returnees	0.76%	3.00%	5.86%	5.57%	5.22%	3.51%
	Attributable fraction, % of diagnoses	13.5	39.1	48.5	42.9	37.1	28.9
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.83%	0.64%	2.19%	3.38%	4.81%	4.59%
	Attributable fraction, % of diagnoses	14.7	8.3	18.1	26.0	34.2	37.8
Military occupation	Background						
	Attributable incidence, % of returnees	4.04%	4.04%	4.04%	4.04%	4.04%	4.04%
	Attributable fraction, % of diagnoses	71.8	52.6	33.4	31.1	28.7	33.3
	Active war service						
	Attributable incidence, % of returnees	0.59%	2.02%	4.40%	4.12%	3.85%	5.18%
	Attributable fraction, % of diagnoses	12.8	34.7	48.0	43.8	35.9	56.8
Combat-specific	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.09%	0.00%	0.82%	1.34%	2.93%	0.00%
	Attributable fraction, % of diagnoses	1.9	0.0	9.0	14.3	27.3	0.0
	Background						
	Attributable incidence, % of returnees	3.94%	3.94%	3.94%	3.94%	3.94%	3.94%
	Attributable fraction, % of diagnoses	85.3	67.7	43.0	41.9	36.8	43.2
Health care	Active war service						
	Attributable incidence, % of returnees	0.85%	3.13%	7.15%	6.18%	6.49%	8.98%
	Attributable fraction, % of diagnoses	14.7	34.7	51.6	41.2	40.8	67.8
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.65%	1.63%	2.43%	4.54%	5.13%	0.00%
	Attributable fraction, % of diagnoses	11.3	18.1	17.5	30.3	32.3	0.0
Other/unspecified	Background						
	Attributable incidence, % of returnees	4.27%	4.27%	4.27%	4.27%	4.27%	4.27%
	Attributable fraction, % of diagnoses	74.0	47.3	30.8	28.5	26.9	32.2
	Active war service						
	Attributable incidence, % of returnees	0.64%	2.21%	4.08%	4.15%	4.07%	2.93%
	Attributable fraction, % of diagnoses	13.0	37.1	44.4	40.6	38.1	31.2
Health care	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.65%	0.10%	1.46%	2.42%	2.96%	2.81%
	Attributable fraction, % of diagnoses	13.2	1.7	15.9	23.7	27.7	29.9
	Background						
	Attributable incidence, % of returnees	3.65%	3.65%	3.65%	3.65%	3.65%	3.65%
	Attributable fraction, % of diagnoses	73.9	61.2	39.7	35.7	34.2	38.9
Other/unspecified	Active war service						
	Attributable incidence, % of returnees	0.60%	1.76%	3.83%	3.75%	3.55%	3.17%
	Attributable fraction, % of diagnoses	11.3	26.9	38.5	34.0	30.3	30.9
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.42%	0.50%	1.83%	3.00%	3.88%	2.79%
	Attributable fraction, % of diagnoses	7.9	7.6	18.4	27.2	33.1	27.2
Background	Attributable incidence, % of returnees	4.29%	4.29%	4.29%	4.29%	4.29%	4.29%
	Attributable fraction, % of diagnoses	80.8	65.5	43.1	38.9	36.6	41.9

TABLE 4b. Attributable incidence rates and attributable fractions of post-traumatic stress disorder (PTSD) diagnoses within 24 months after return from Iraq/Afghanistan, by determinants of likelihood of diagnosis, active component, U.S. Armed Forces, by follow-up period, April 2003–December 2013

b. PTSD

Determinants of the likelihood of diagnosis		2003–2005	2005–2007	2007–2009	2009–2011	2011–2012	2012–2013
Overall	Active war service						
	Attributable incidence, % of returnees	2.48%	4.31%	4.44%	3.95%	4.29%	2.38%
	Attributable fraction, % of diagnoses	82.1	81.5	73.9	66.1	63.8	46.3
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.26%	0.70%	1.29%	1.75%	2.15%	2.48%
	Attributable fraction, % of diagnoses	8.6	13.2	21.4	29.2	31.9	48.2
Background	Attributable incidence, % of returnees	0.28%	0.28%	0.28%	0.28%	0.28%	0.28%
	Attributable fraction, % of diagnoses	9.4	5.3	4.7	4.7	4.2	5.5
Service Army	Active war service						
	Attributable incidence, % of returnees	3.76%	5.93%	6.48%	4.45%	4.93%	2.42%
	Attributable fraction, % of diagnoses	83.6	79.2	71.3	55.8	56.3	35.2
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.44%	1.26%	2.31%	3.23%	3.52%	4.16%
	Attributable fraction, % of diagnoses	9.7	16.8	25.4	40.5	40.3	60.5
Background	Attributable incidence, % of returnees	0.30%	0.30%	0.30%	0.30%	0.30%	0.30%
	Attributable fraction, % of diagnoses	6.6	4.0	3.3	3.7	3.4	4.3
Navy	Active war service						
	Attributable incidence, % of returnees	1.69%	4.34%	2.46%	2.90%	3.59%	4.39%
	Attributable fraction, % of diagnoses	100.0	100.0	77.0	73.4	76.1	100.0
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.00%	0.00%	0.73%	1.05%	1.13%	0.00%
	Attributable fraction, % of diagnoses	0.0	0.0	23.0	26.6	23.9	0.0
Background	Attributable incidence, % of returnees	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Attributable fraction, % of diagnoses	0.0	0.0	0.0	0.0	0.0	0.0
Air Force	Active war service						
	Attributable incidence, % of returnees	0.54%	0.98%	1.37%	1.21%	1.71%	1.80%
	Attributable fraction, % of diagnoses	70.1	81.1	76.3	60.2	68.0	88.7
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.00%	0.00%	0.20%	0.57%	0.57%	0.00%
	Attributable fraction, % of diagnoses	0.0	0.0	10.9	28.4	22.8	0.0
Background	Attributable incidence, % of returnees	0.23%	0.23%	0.23%	0.23%	0.23%	0.23%
	Attributable fraction, % of diagnoses	29.9	18.9	12.8	11.4	9.1	11.3
Marine Corps	Active war service						
	Attributable incidence, % of returnees	2.46%	4.67%	3.53%	5.72%	4.48%	3.96%
	Attributable fraction, % of diagnoses	81.8	80.8	66.5	94.1	69.8	91.7
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.19%	0.75%	1.42%	0.00%	1.58%	0.00%
	Attributable fraction, % of diagnoses	6.3	13.0	26.8	0.0	24.6	0.0
Background	Attributable incidence, % of returnees	0.36%	0.36%	0.36%	0.36%	0.36%	0.36%
	Attributable fraction, % of diagnoses	11.9	6.2	6.7	5.9	5.6	8.3
Sex Male	Active war service						
	Attributable incidence, % of returnees	2.57%	4.62%	4.69%	4.20%	4.48%	2.67%
	Attributable fraction, % of diagnoses	84.9	84.4	76.6	69.7	66.5	52.6
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.25%	0.65%	1.22%	1.62%	2.05%	2.20%
	Attributable fraction, % of diagnoses	8.3	11.8	20.0	26.9	30.5	43.4
Background	Attributable incidence, % of returnees	0.21%	0.21%	0.21%	0.21%	0.21%	0.21%
	Attributable fraction, % of diagnoses	6.8	3.8	3.4	3.4	3.1	4.1
Female	Active war service						
	Attributable incidence, % of returnees	2.25%	2.21%	2.88%	2.49%	3.37%	4.93%
	Attributable fraction, % of diagnoses	75.5	58.3	55.9	44.5	50.7	87.1
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.00%	0.85%	1.54%	2.38%	2.55%	0.00%
	Attributable fraction, % of diagnoses	0.0	22.4	29.9	42.5	38.3	0.0
Background	Attributable incidence, % of returnees	0.73%	0.73%	0.73%	0.73%	0.73%	0.73%
	Attributable fraction, % of diagnoses	24.5	19.3	14.2	13.0	11.0	12.9
Age group <25 years	Active war service						
	Attributable incidence, % of returnees	2.87%	5.53%	5.35%	4.72%	4.89%	4.33%
	Attributable fraction, % of diagnoses	81.0	83.7	77.3	72.5	72.3	92.4
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.32%	0.72%	1.21%	1.43%	1.51%	0.00%
	Attributable fraction, % of diagnoses	9.0	10.9	17.5	22.0	22.4	0.0
Background	Attributable incidence, % of returnees	0.36%	0.36%	0.36%	0.36%	0.36%	0.36%
	Attributable fraction, % of diagnoses	10.1	5.4	5.2	5.5	5.3	7.6

Table 4b (cont.)

Determinants of the likelihood of diagnosis		2003–2005	2005–2007	2007–2009	2009–2011	2011–2012	2012–2013
25–35 years	Active war service						
	Attributable incidence, % of returnees	2.36%	3.85%	4.09%	3.62%	4.32%	2.35%
	Attributable fraction, % of diagnoses	83.6	80.6	74.4	64.3	65.5	46.9
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.24%	0.70%	1.19%	1.79%	2.06%	2.44%
	Attributable fraction, % of diagnoses	8.5	14.7	21.6	31.8	31.2	48.7
>35 years	Background						
	Attributable incidence, % of returnees	0.22%	0.22%	0.22%	0.22%	0.22%	0.22%
	Attributable fraction, % of diagnoses	7.8	4.6	4.0	3.9	3.3	4.4
	Active war service						
	Attributable incidence, % of returnees	2.13%	2.97%	3.47%	3.39%	3.18%	6.37%
	Attributable fraction, % of diagnoses	100.0	76.6	65.1	57.2	45.3	100.0
Race/ethnicity	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.00%	0.90%	1.86%	2.54%	3.84%	0.00%
	Attributable fraction, % of diagnoses	0.0	23.4	34.9	42.8	54.7	0.0
	Background						
	Attributable incidence, % of returnees	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
	Attributable fraction, % of diagnoses	0.0	0.0	0.0	0.0	0.0	0.0
White, non-Hispanic	Active war service						
	Attributable incidence, % of returnees	2.55%	4.53%	4.60%	4.27%	4.50%	2.47%
	Attributable fraction, % of diagnoses	80.7	81.5	74.1	70.3	68.0	50.2
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.31%	0.73%	1.31%	1.50%	1.82%	2.15%
	Attributable fraction, % of diagnoses	9.8	13.1	21.1	24.7	27.5	43.7
Black, non-Hispanic	Background						
	Attributable incidence, % of returnees	0.30%	0.30%	0.30%	0.30%	0.30%	0.30%
	Attributable fraction, % of diagnoses	9.5	5.4	4.8	4.9	4.5	6.1
	Active war service						
	Attributable incidence, % of returnees	2.1%	3.1%	3.6%	3.2%	3.4%	2.2%
	Attributable fraction, % of diagnoses	82.5	77.8	68.0	55.6	50.7	38.2
Hispanic	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.2%	0.6%	1.4%	2.3%	3.1%	3.3%
	Attributable fraction, % of diagnoses	6.8	15.6	26.9	39.8	45.3	57.1
	Background						
	Attributable incidence, % of returnees	0.269%	0.269%	0.269%	0.269%	0.269%	0.269%
	Attributable fraction, % of diagnoses	10.8	6.6	5.1	4.6	4.0	4.7
Other/unspecified	Active war service						
	Attributable incidence, % of returnees	2.7%	6.1%	5.1%	4.1%	5.4%	5.7%
	Attributable fraction, % of diagnoses	81.8	96.5	77.2	60.4	68.1	96.2
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.4%	0.0%	1.3%	2.4%	2.3%	0.0%
	Attributable fraction, % of diagnoses	11.6	0.0	19.4	36.3	29.1	0.0
Military occupation	Background						
	Attributable incidence, % of returnees	0.221%	0.221%	0.221%	0.221%	0.221%	0.221%
	Attributable fraction, % of diagnoses	6.7	3.5	3.3	3.3	2.8	3.8
	Active war service						
	Attributable incidence, % of returnees	2.8%	3.6%	3.9%	3.2%	3.9%	4.7%
	Attributable fraction, % of diagnoses	100.0	80.8	76.7	68.0	65.3	100.0
Combat-specific	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.0%	0.0%	1.2%	1.5%	2.1%	0.0%
	Attributable fraction, % of diagnoses	0.0	0.0	23.3	32.0	34.7	0.0
	Background						
	Attributable incidence, % of returnees	0.000%	0.000%	0.000%	0.000%	0.000%	0.000%
	Attributable fraction, % of diagnoses	0.0	0.0	0.0	0.0	0.0	0.0
Health care	Active war service						
	Attributable incidence, % of returnees	3.26%	6.22%	5.81%	5.21%	4.97%	5.21%
	Attributable fraction, % of diagnoses	81.8	85.4	74.6	69.6	62.6	95.9
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.51%	0.84%	1.76%	2.06%	2.75%	0.00%
	Attributable fraction, % of diagnoses	12.7	11.5	22.6	27.5	34.6	0.0
Other/unspecified	Background						
	Attributable incidence, % of returnees	0.22%	0.22%	0.22%	0.22%	0.22%	0.22%
	Attributable fraction, % of diagnoses	5.5	3.0	2.8	2.9	2.8	4.1
	Active war service						
	Attributable incidence, % of returnees	5.34%	6.95%	7.22%	6.06%	6.82%	7.33%
	Attributable fraction, % of diagnoses	92.4	83.4	81.5	75.7	71.6	94.3
Health care	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.00%	0.94%	1.20%	1.51%	2.26%	0.00%
	Attributable fraction, % of diagnoses	0.0	11.3	13.5	18.8	23.7	0.0
	Background						
	Attributable incidence, % of returnees	0.44%	0.44%	0.44%	0.44%	0.44%	0.44%
	Attributable fraction, % of diagnoses	7.6	5.3	5.0	5.5	4.6	5.7
Other/unspecified	Active war service						
	Attributable incidence, % of returnees	1.87%	3.23%	3.41%	3.18%	3.60%	2.00%
	Attributable fraction, % of diagnoses	79.3	77.7	69.5	61.3	60.9	42.2
	Enhanced ascertainment						
	Attributable incidence, % of returnees	0.21%	0.64%	1.22%	1.73%	2.03%	2.46%
	Attributable fraction, % of diagnoses	8.7	15.5	24.8	33.3	34.3	51.8
Background	Attributable incidence, % of returnees	0.28%	0.28%	0.28%	0.28%	0.28%	0.28%
	Attributable fraction, % of diagnoses	12.0	6.8	5.8	5.5	4.8	6.0