

Deployment Health Assessments

U.S. Armed Forces

April 2016

(Data through March 2016)

Armed Forces Health Surveillance Branch

Update: Deployment Health Assessments, U.S. Armed Forces, April 2016

Since January 2003, peaks and troughs in the monthly numbers of pre- and post-deployment health assessment forms transmitted to the Armed Forces Health Surveillance Branch generally corresponded to times of departure and return of large numbers of service members who deployed. Between April 2006 and March 2016, the number of post-deployment reassessment (PDHRA) forms per month ranged from 15,114 to 45,768 (Figure 1). For the most recent 12 months (April 2015–March 2016), the average monthly numbers of pre- and post-deployment assessments (18,269 and 10,907, respectively) were the second lowest and lowest of any 12-month period since the first such period ended in January 2004 (Table 1).

During the past 12 months, the proportions of returned deployers who rated their health as either “fair” or “poor” were 5.3%–7.8% on post-deployment health assessment questionnaires and 7.1%–10.1% on PDHRA questionnaires (Figure 2). The proportions of service members who rated their health as either “fair” or “poor” on the pre-deployment questionnaires ranged from 2.9% to 4.5% during the past 12 months.

In general, the proportions of reserve component members reporting “exposure concerns” and “health concerns” on post-deployment assessments and reassessments have been consistently greater than among active component members (Table 2, Figure 3). Service members in the active component of the Marine Corps and reserve component of the Army were more likely to report exposure concerns 3–6 months after, compared to the time of, return from deployment (Table 2).

At the time of return from deployment, service members in the active and reserve component of the Army, active component of the Navy, and reserve component of the Marine Corps were the most likely of all deployers to receive mental health referrals; 3–6 months after returning, at the time of the PDHRA, reservists in all services were more likely than their active component counterparts to receive mental health referrals (Table 2).

Table 1. Numbers of deployment-related health assessment forms, by month, U.S. Armed Forces, April 2015–March 2016

	Pre-deployment assessment DD 2795		Post-deployment assessment DD 2796		Post-deployment reassessment DD 2900	
	No.	%	No.	%	No.	%
Total	219,225	100	130,878	100	243,669	100
2015						
April	20,833	9.5	10,380	7.9	22,611	9.3
May	18,922	8.6	7,373	5.6	20,097	8.2
June	17,933	8.2	10,382	7.9	20,258	8.3
July	17,882	8.2	13,067	10.0	19,771	8.1
August	19,842	9.1	10,456	8.0	21,384	8.8
September	18,503	8.4	14,120	10.8	17,149	7.0
October	19,484	8.9	13,052	10.0	18,337	7.5
November	12,654	5.8	10,293	7.9	19,125	7.8
December	16,413	7.5	11,750	9.0	19,415	8.0
2016						
January	20,599	9.4	9,508	7.3	21,742	8.9
February	18,080	8.2	7,538	5.8	20,974	8.6
March	18,080	8.2	12,959	9.9	22,806	9.4

Source: Defense Medical Surveillance System (DMSS) as of 07APR2016
Prepared by Armed Forces Health Surveillance Branch (AFHSB)

Figure 2. Proportion of deployment health assessment forms with self-assessed health status as “fair” or “poor,” by month, U.S. Armed Forces, April 2015–March 2016





Figure 1. Numbers of deployment health assessment and reassessment forms, by month, U.S. Armed Forces, January 2003–March 2016



Data Source: Defense Medical Surveillance System

Table 2. Percentage of service members who endorsed selected questions or received referrals on health assessment forms, U.S. Armed Forces, April 2015–March 2016

	Army			Navy			Air Force			Marine Corps			All service members		
	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900
Active component	n= 94,443	n= 38,694	n= 88,805	n= 6,730	n= 6,137	n= 8,732	n= 37,435	n= 34,901	n= 69,882	n= 17,654	n= 13,691	n= 13,985	n= 156,262	n= 93,423	n= 181,404
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
General health "fair" or "poor"	5.8	8.9	4.8	1.9	5.0	3.5	1.2	2.7	1.2	3.0	5.8	6.9	4.2	5.8	3.5
Health concerns, not wound or injury	7.4	20.2	6.8	3.2	10.8	7.1	3.1	7.4	7.5	3.4	8.7	8.2	5.7	13.1	7.2
Health worse now than before deployed	na	72.0	32.8	na	70.0	54.3	na	60.6	27.4	na	71.1	66.3	na	67.5	34.3
Exposure concerns	na	12.7	5.5	na	14.2	10.4	na	11.0	2.9	na	9.3	15.3	na	11.6	5.5
PTSD symptoms (2 or more)	na	6.1	3.7	na	3.5	7.2	na	2.1	3.3	na	2.8	7.5	na	4.0	4.0
Depression symptoms (any)	na	21.6	9.7	na	15.3	17.1	na	8.3	9.4	na	14.6	19.6	na	15.2	10.7
Referral indicated by provider (any)	7.4	19.2	17.6	5.0	28.6	12.8	5.8	12.5	8.8	5.3	18.1	14.4	6.7	17.2	13.7
Mental health referral indicated ^a	2.1	3.2	14.4	0.7	4.1	10.3	0.9	1.5	7.3	0.6	2.0	12.4	1.6	2.4	11.3
Medical visit following referral ^b	97.4	93.3	94.4	84.3	47.8	76.3	97.4	79.7	96.6	87.1	52.8	82.3	96.4	76.8	93.0
	Army			Navy			Air Force			Marine Corps			All service members		
	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900	Pre-deploy DD2795	Post-deploy DD2796	Reassessmt DD2900
Reserve component	n= 21,300	n= 16,775	n= 14,516	n= 2,276	n= 1,861	n= 3,406	n= 13,318	n= 10,174	n= 21,044	n= 594	n= 240	n= 8,863	n= 37,488	n= 29,050	n= 47,829
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
General health "fair" or "poor"	1.3	10.4	13.3	0.2	8.9	5.7	0.9	4.5	1.5	0.8	9.6	0.7	1.1	8.2	5.2
Health concerns, not wound or injury	2.8	22.9	19.6	1.1	40.5	8.5	1.7	9.5	3.6	2.0	27.9	2.6	2.3	19.4	8.6
Health worse now than before deployed	na	69.4	77.5	na	76.1	43.5	na	66.7	25.0	na	74.6	5.1	na	68.9	38.6
Exposure concerns	na	17.3	21.9	na	37.7	18.1	na	16.3	3.5	na	22.1	1.4	na	18.3	9.7
PTSD symptoms (2 or more)	na	7.4	13.8	na	6.4	11.7	na	1.6	2.6	na	8.3	7.4	na	5.3	7.5
Depression symptoms (any)	na	20.2	23.4	na	17.8	22.8	na	8.1	7.2	na	17.9	19.2	na	15.8	15.5
Referral indicated by provider (any)	4.4	28.4	45.1	4.2	40.7	25.8	3.0	15.4	8.4	4.5	55.8	14.1	3.9	24.9	21.9
Mental health referral indicated ^a	0.9	4.2	44.6	0.3	2.9	25.4	0.4	1.2	8.1	1.5	4.6	12.8	0.7	3.0	21.3
Medical visit following referral ^b	90.5	98.1	37.5	88.5	94.7	44.4	62.5	53.4	45.1	100.0	49.1	37.0	83.0	87.3	39.1

^aIncludes behavioral health, combat stress and substance abuse referrals.

^bRecord of inpatient or outpatient visit within 6 months after referral.

Figure 3. Proportion of service members who endorsed exposure concerns on post-deployment health assessments or reassessments, by component, U.S. Armed Forces, January 2003–March 2016

