## Deployment Health Assessments U.S. Armed Forces December 2019

(Data through November 2019)

Armed Forces Health Surveillance Branch

Report Date: 19 December 2019

## Update: Deployment Health Assessments, U.S. Armed Forces, December 2019

Since January 2003, pronounced fluctuations in the monthly numbers of pre- and post-deployment health assessment (PDHA) forms transmitted to the Armed Forces Health Surveillance Branch (AFHSB) generally have corresponded to times of departure and return of large numbers of service members who deployed. From October 2014 through November 2019, monthly counts of pre-deployment health assessments showed the greatest variation compared to the other deployment health assessment forms (Figure 1). Since June 2017, monthly counts of post deployment health reassessment (PDHRA) forms have been consistently lower than the monthly PDHA counts.


The average monthly number of PDHRA forms (7,491) during the most recent 12 months (December 2018–November 2019) was the single lowest 1-year average since July 2006 (**Table 1**, **data not shown**). The average monthly number of pre-deployment health assessment forms (19,546) during this period was the lowest 1-year average since July 2017 (**Table 1**, **data not shown**). For the purposes of this report about service members' responses to the assessment questions, it is important to note that the denominator is the total number of service members who completed deployment-related assessments, not the total number of service members eligible for the assessments.

During the past 12 months, the percentages of total returned deployers who rated their health as either "fair" or "poor" ranged from 7.2% to 10.1% on PDHA questionnaires and from 7.0% to 8.2% on PDHRA questionnaires (Figure 2). The percentages of total service members who rated their health as either "fair" or "poor" on pre-deployment questionnaires during this period varied between 3.4% and 4.8%.

In general, the percentages of "exposure concerns" self-reported on PDHA and PDHRA forms were higher among reserve component members than among active component members; a similar pattern is apparent for the percentages of "health concerns" self-reported on PDHA forms (Table 2, Figure 3). Active and reserve component Marine Corps members were 1.8 and 1.5 times as likely, respectively, to report 2 or more PTSD symptoms at the time of the PDHRA compared to the time of the PDHA (Table 2).

At the time of return from deployment, 17.6% of active component and reserve component service members reported any depression symptoms. Service members in the active and reserve components were 4.1 and 6.2 times as likely, respectively, to receive a mental health referral 3-6 months after deployment, compared to the time of their return from deployment. Moreover, at the time of the PDHRA, reservists were 1.8 times as likely to receive a mental health referral as members of the active component (**Table 2**).

Figure 1. Numbers of deployment health assessment and reassessment forms, by month, U.S. Armed Forces, January 2003–November 2019




**Table 1.** Number of deployment-related health assessment forms, by month, U.S. Armed Forces, December 2018–November 2019

	Pre-depl assess DD 2	sment	Post-dep assess DD 2	ment	Post-deployment reassessment DD 2900			
	No.	%	No.	%	No.	%		
Total	234,551	100.0	166,869	100.0	89,892	100.0		
Average	19,546		13,906		7,491			
2018								
December	16,558	7.1	11,612	7.0	5,892	6.6		
2019								
January	28,609	12.2	16,356	9.8	9,653	10.7		
February	19,227	8.2	14,540	8.7	9,324	10.4		
March	17,398	7.4	17,424	10.4	6,794	7.6		
April	19,825	8.5	11,649	7.0	6,847	7.6		
May	21,242	9.1	12,011	7.2	9,025	10.0		
June	19,438	8.3	13,189	7.9	8,018	8.9		
July	20,375	8.7	15,845	9.5	8,022	8.9		
August	26,164	11.2	9,849	5.9	7,670	8.5		
September	14,987	6.4	18,158	10.9	5,032	5.6		
October	17,304	7.4	15,087	9.0	6,692	7.4		
November	13,424	5.7	11,149	6.7	6,923	7.7		

Source: Defense Medical Surveillance System (DMSS) as of 09-DEC-2019 Prepared by Armed Forces Health Surveillance Branch (AFHSB)

**Figure 2.** Percentages of deployment health assessment forms with self-assessed health status rated as either "fair" or "poor," by month, U.S. Armed Forces. December 2018–November 2019




**Table 2.** Percentage of service members who endorsed selected questions or received referrals on health assessment forms, U.S. Armed Forces, December 2018–November 2019

	Army			Navy		Air Force		Marine Corps			All active component service members				
	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessn DD2900
Active component	n= 91,576 %	n= 55,941 %	n= 24,933 %	n= 5,566 %	n= 5,522 %	n= 4,478 %	n= 38,530 %	n= 32,022 %	n= 28,386 %	n= 17,976 %	n= 18,358 %	n= 10,528 %	n= 153,648 %	n= 111,843 %	n= 68,325 %
General health "fair" or "poor"	6.2	10.5	10.8	2.4	6.3	6.1	1.6	3.9	4.0	3.3	7.5	9.2	4.6	7.9	7.4
Health concerns, not wound or injury	6.3	17.9	11.5	4.7	13.8	10.7	3.2	10.8	4.0 8.1	4.6	10.9	10.7	5.2	7.9 14.5	9.9
Health worse now than before deployed	na	72.7	73.5	na	69.2	76.2	na	63.2	70.7	na	71.4	78.3	na	69.6	73.3
Exposure concerns	na	13.3	12.0	na	22.3	24.0	na	14.1	9.6	na	10.5	14.6	na	13.5	12.2
PTSD symptoms (two or more)	na	6.2	8.2	na	4.5	6.6	na	2.6	3.1	na	3.2	5.8	na	4.6	5.6
Depression symptoms (any)	na	21.4	21.3	na	19.6	20.5	na	10.1	10.1	na	18.1	20.1	na	17.6	16.4
Referral indicated by provider (any)	6.2	14.8	11.3	7.1	25.7	19.9	7.7	14.9	9.5	6.0	21.4	16.9	6.6	16.4	12.0
Mental health referral indicated <sup>a</sup>	1.5	2.4	9.0	1.4	4.9	17.2	1.2	2.2	8.3	1.0	2.4	15.0	1.4	2.5	10.2
Medical visit following referral <sup>b</sup>	92.5	96.4	97.6	82.3	86.4	80.5	97.8	98.3	97.5	81.6	93.6	94.2	92.1	95.6	94.8
	Army		Navy		Air Force		Marine Corps			All reserve component service members					
	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessn DD2900
Reserve component	n= 37,666 %	n= 30,738 %	n= 6,389 %	n= 2,602 %	n= 2,433 %	n= 2,367 %	n= 12,912 %	n= 13,669 %	n= 11,537 %	n= 1,864 %	n= 1,981 %	n= 572 %	n= 55,044 %	n= 48,821 %	n= 20,865 %
Company								5.9						9.3	7.3
General health "fair" or "poor"  Health concerns, not wound or injury	1.7 2.9	10.5 19.6	11.7 11.8	0.8 3.2	11.0 38.0	9.2 12.0	1.0 1.7	5.9 12.6	4.5 6.2	1.1 2.3	10.8 19.0	8.2 7.2	1.5 2.6	9.3 18.5	7.3 8.6
Health worse now than before deployed	na	69.6	74.0	na	73.4	76.2	na	67.2	71.8	na na	74.3	79.0	na	69.3	73.2
Exposure concerns	na	18.4	18.2	na	43.6	37.3	na	19.9	10.9	na	14.1	13.5	na	19.9	16.2
PTSD symptoms (two or more)	na	7.6	9.6	na	7.4	9.6	na	2.6	2.8	na	5.6	8.4	na	6.1	5.8
Depression symptoms (any)	na	20.7	19.8	na	20.2	22.8	na	10.1	8.1	na	17.6	18.4	na	17.6	13.6
Referral indicated by provider (any)	3.5	21.2	16.6	5.3	29.6	34.2	4.2	15.9	18.9	8.2	27.3	30.2	3.9	20.4	20.3
Mental health referral indicated <sup>a</sup>	0.8	3.3	11.7	0.8	3.4	33.5	0.5	1.9	18.4	1.9	5.2	29.0	0.7	3.0	18.4
Medical visit following referral <sup>b</sup> PTSD. post-traumatic stress disorder	87.6	99.1	49.7	91.9	97.6	62.5	81.4	84.2	81.9	87.4	85.4	49.2	85.9	95.5	68.1

PTSD, post-traumatic stress disorder

**Figure 3.** Percentages of service members with completed assessments who endorsed exposure concerns on post-deployment health assessments or reassessments, by component, by month, U.S. Armed Forces, January 2003–November 2019



<sup>&</sup>lt;sup>a</sup>Includes behavioral health, combat stress and substance abuse referrals.

<sup>&</sup>lt;sup>b</sup>Record of inpatient or outpatient visit within 6 months after referral.