Deployment Health Assessments

U.S. Armed Forces

October 2020

(Data through September 2020)

Armed Forces Health Surveillance Branch

Update: Deployment Health Assessments, U.S. Armed Forces, October 2020

Since January 2003, pronounced fluctuations in the monthly numbers of pre- and post-deployment health assessment (PDHA) forms transmitted to the Armed Forces Health Surveillance Branch (AFHSB) generally have corresponded to times of departure and return of large numbers of service members who deployed. For the purposes of this report about service members' responses to the assessment questions, it is important to note that the denominator is the total number of service members who completed deployment-related assessments, not the total number of service members eligible for the assessments.


From January 2015 through September 2020, monthly counts of pre-deployment health assessment forms showed the greatest variation compared to the other deployment health assessment forms (Figure 1). The most pronounced fluctuations in monthly counts of pre-deployment health assessment forms during this period occurred from June 2017 through September 2020 (Figure 1). Moreover, the average monthly number of pre-deployment health assessment forms (22,511) during the most recent 12 months was the highest 1-year average since February 2019 (Table 1, data not shown). For the past 8 years, counts of post-deployment health reassessment (PDHRA) forms showed a pronounced downward trend; since June 2017, the monthly counts of these questionnaires were consistently lower than the monthly counts of PDHA forms.

During the past 12 months, the percentages of total returned deployers who rated their health as either "fair" or "poor" ranged from 5.6% to 10.0% on PDHA questionnaires and from 6.4% to 9.3% on PDHRA questionnaires (**Figure 2**). The percentages of total service members who rated their health as either "fair" or "poor" on pre-deployment questionnaires during this period varied between 2.2% and 5.6%.

In general, the percentages of "exposure concerns" self-reported on PDHA and PĎHRA forms were higher among reserve component members than among active component members; a similar pattern is apparent for the percentages of "health concerns" self-reported on PDHA forms (**Table 2**, **Figure 3**). Active component Marine Corps members were 1.6 times as likely to report 2 or more PTSD symptoms at the time of the PDHRA compared to the time of the PDHA (**Table 2**).

At the time of return from deployment, 18.9% of active component and 15.5% of reserve component service members reported any depression symptoms. Service members in the active and reserve components were 3.6 and 8.6 times as likely, respectively, to receive a mental health referral 3-6 months after deployment, compared to the time of their return from deployment. Moreover, at the time of the PDHRA, reservists were 1.9 times as likely to receive a mental health referral as members of the active component (Table 2).

Figure 1. Numbers of deployment health assessment and reassessment forms, a by month, U.S. Armed Forces, January 2003–September 2020



No., number


^aPre-deployment health assessment forms are completed 30–90 days prior to departure for deployment. Post-deployment assessment forms are completed 1–30 days after return from deployment. Post-deployment re-assessment forms are completed 90-180 days after return from deployment.

Table 1. Number of deployment-related health assessment forms, by month, U.S. Armed Forces, October 2019–September 2020

	Pre-depl assess DD 2	ment	Post-dep assess DD 2	ment	Post-deployment reassessment DD 2900			
	No.	%	No.	%	No.	%		
Total	270,128	100.0	182,526	100.0	87,046	100.0		
Average	22,511		15,211		7,254			
2019								
October	17,519	6.5	15,728	8.6	6,967	8.0		
November	14,129	5.2	11,630	6.4	7,829	9.0		
December	20,558	7.6	10,374	5.7	6,926	8.0		
2020								
January	40,195	14.9	11,497	6.3	9,245	10.6		
February	21,107	7.8	11,971	6.6	9,224	10.6		
March	22,016	8.2	11,345	6.2	7,489	8.6		
April	23,610	8.7	8,181	4.5	5,448	6.3		
May	26,302	9.7	21,739	11.9	6,650	7.6		
June	19,586	7.3	25,476	14.0	6,739	7.7		
July	23,520	8.7	20,915	11.5	5,910	6.8		
August	20,593	7.6	16,140	8.8	5,836	6.7		
September	20,993	7.8	17,530	9.6	8,783	10.1		
No., number.								

Source: Defense Medical Surveillance System (DMSS) as of 08-OCT-2020 Prepared by Armed Forces Health Surveillance Branch (AFHSB)

Figure 2. Percentages of deployment health assessment forms with self-assessed health status rated as either "fair" or "poor," by month, U.S. Armed Forces, October 2019–September 2020




Data Source: Defense Medical Surveillance System

Table 2. Percentage of service members who endorsed selected questions or received referrals on health assessment forms, U.S. Armed Forces, October 2019-September 2020

		Army		Navy		Air Force		Marine Corps			All active component service members				
	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessm DD2900
Active component	n= 99,942	n= 53,135	n= 23,396	n= 6,123	n= 5,469	n= 5,504	n= 40,381	n= 32,778	n= 26,408	n= 16,308	n= 16,107	n= 10,680	n= 162,754	n= 107,489	n= 65,988
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
General health "fair" or "poor"	6.8	10.5	10.5	2.8	7.3	7.1	1.7	4.6	4.4	3.8	7.0	9.4	5.1	8.0	7.6
Health concerns, not wound or injury	7.1	18.1	10.7	5.6	15.7	12.2	3.7	10.7	9.1	4.6	11.8	10.9	5.9	14.8	10.2
Health worse now than before deployed	na	72.7	73.8	na	73.4	76.3	na	64.0	71.7	na	71.3	76.3	na	69.9	73.6
Exposure concerns	na	12.8	12.6	na	16.9	17.4	na	14.4	10.5	na	8.8	11.6	na	12.9	12.0
PTSD symptoms (two or more)	na	7.0	8.7	na	7.3	7.8	na	3.9	3.6	na	3.3	5.2	na	5.5	6.0
Depression symptoms (any)	na	22.6	22.2	na	23.2	24.0	na	12.7	10.9	na	17.8	19.6	na	18.9	17.4
Referral indicated by provider (any)	5.2	12.7	12.5	6.3	28.8	18.7	7.4	17.8	9.9	6.9	20.1	16.0	6.0	16.2	12.5
Mental health referral indicateda	1.5	2.6	9.9	1.0	6.7	16.4	1.5	3.4	8.8	0.8	2.3	14.5	1.4	3.0	10.8
Medical visit following referral ^b	98.1	97.9	97.2	87.2	81.7	87.8	99.2	98.2	99.1	95.8	89.5	90.6	97.7	94.9	95.1
	Army		Navy		Air Force		Marine Corps			All reserve component service members					
	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessmt DD2900	Pre- deploy DD2795	Post- deploy DD2796	Reassessn DD2900
Reserve component	n= 50,217	n= 5,630	n= 3,354	n= 2,159	n= 2,350	n= 19,588	n= 15,735	n= 11,361	n= 1,098	n= 764	n= 1,018	n= 71,320	n= 68,875	n= 20,359	n= 18,707
	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
General health "fair" or "poor"	1.4	7.8	8.4	1.2	10.6	9.9	0.8	4.5	4.6	0.7	8.2	8.3	1.2	7.1	6.4
Health concerns, not wound or injury	2.4	14.3	10.9	3.2	37.2	13.5	1.8	9.2	6.2	2.4	21.2	8.8	2.3	13.9	8.5
Health worse now than before deployed	na	72.6	70.0	na	75.1	77.8	na	73.4	74.9	na	76.6	76.3	na	72.9	74.0
Exposure concerns	na	15.5	11.7	na	39.6	35.9	na	15.4	11.8	na	20.2	16.2	na	16.3	14.8
PTSD symptoms (two or more)	na	6.4	7.7	na	9.4	10.2	na	2.9	3.0	na	6.7	9.3	na	5.7	5.5
Depression symptoms (any)	na	16.9	16.1	na	23.0	23.9	na	9.8	9.2	na	22.5	19.2	na	15.5	13.3
Referral indicated by provider (any)	2.3	13.8	15.8	7.6	46.7	38.8	3.3	16.7	20.6	7.1	44.1	31.9	2.9	15.8	21.9
Mental health referral indicated ^a	0.5	2.1	11.2	0.8	7.8	38.1	0.5	2.2	20.0	0.9	8.1	31.5	0.5	2.4	20.3
Medical visit following referral ^b PTSD. post-traumatic stress disorder	81.2	99.7	55.9	96.1	90.3	56.4	91.6	82.4	77.7	97.0	98.9	35.4	86.3	93.9	67.3

PTSD, post-traumatic stress disorder

Figure 3. Percentages of service members with completed assessments who endorsed exposure concerns on post-deployment health assessments or reassessments, by component, by month, U.S. Armed Forces, January 2003–September 2020



^aIncludes behavioral health, combat stress and substance abuse referrals.

^bRecord of inpatient or outpatient visit within 6 months after referral.