**TABLE 5.** Adjusted Hazard Ratios for Time of Reporting Nine Most Frequent Reportable Medical Events Among Other Department of War Beneficiaries to DRSi by Period, Estimated Medical Personnel Category and Event Category, January 1, 2016–November 1, 2023

| Variable | Nine Most Frequent RMEs[a] | | | Chlamydia, COVID-19 Exclusion[b] | | |
|---|---|---|---|---|---|---|
| | Hazard Ratio | 95% CI | | Hazard Ratio | 95% CI | |
| | | LL | UL | | LL | UL |
| Period | | | | | | |
| Post-transition vs. pre-transition—Reference | 0.64 | 0.62 | 0.67 | 0.85 | 0.79 | 0.92 |
| Estimated medical personnel category[c] | | | | | | |
| Small vs. large—Reference | 1.12 | 1.07 | 1.17 | 1.18 | 1.05 | 1.31 |
| Medium vs. large—Reference | 0.96 | 0.94 | 0.99 | 1.00 | 0.93 | 1.07 |
| RME category | | | | | | |
| Gastrointestinal infection vs. STI—Reference | 0.52 | 0.49 | 0.54 | 0.92 | 0.87 | 0.96 |
| Rabies prophylaxis vs. STI—Reference | 0.45 | 0.41 | 0.49 | 0.78 | 0.71 | 0.86 |
| Respiratory infection vs. STI—Reference | 0.34 | 0.33 | 0.35 | 0.23 | 0.21 | 0.25 |

Abbreviations: DRSi, Disease Reporting System internet; RMEs, reportable medical events; COVID-19, coronavirus disease 2019; CI: confidence interval; LL, lower limit; upper limit; vs., versus; STI, sexually transmitted infection.

[a] 30,116 censored (42.0%) of 71,700 records.

[b] 11,917 censored (59.8%) of 19,940 records.

[c] Estimated medical personnel category full-time equivalent person-months: small, 1,413–3,314; medium, 3,315–8,401; large, 8,402–90,242.